


# EOS



The VCP News, page 722

Eos, Transactions, American Geophysical Union

Vol. 65, No. 39, Pages 721-728

September 25, 1984

New phone numbers (will be published in Membership Directory)

Office ( ) Home ( )

Please print or type new address

American Geophysical Union  
2000 Florida Avenue, N.W.  
Washington, DC 20009Or, for faster action, call toll free  
800-424-2438; in the Washington,  
DC, area, call 462-6993.

## Tectonophysics

1950 Tectonophysics (Plate tectonics) has been a critical component of the Earth's evolution. It is the study of the forces that drive the movement of the Earth's plates and the resulting geological features.

Active fold-and-thrust belts or subduction accretionary complexes can be modeled as critically tapered wedges of material on the verge of Coulomb failure everywhere, overlying a basal décollement where frictional sliding is occurring. Ignoring cohesion, the four strength parameters needed to describe a critical Coulomb wedge are its internal and basal coefficients of friction  $\mu$  and  $\mu_b$  and its internal and basal fluid-grain pressure ratios  $\lambda$  and  $\lambda_b$ . An exact relation between the surface slope  $\alpha$  and basal dip  $\beta$  of a non-cohesive critical wedge with uniform properties is derived. The state of stress within such a wedge has the same orientation everywhere, and is constant if  $\beta$  is and  $\mu$  and  $\lambda$  are constant. A coefficient of internal friction  $\mu = 1$  is consistent with the known surface slope, basal dip and pore-fluid pressures in the active fold-and-thrust belt of western Taiwan, assuming Byerlee's law  $\tau = \mu \sigma_n$  is valid on the base. The wide variety of tectonic styles observed to occur along convergent margins, including subduction arcs, active accretion, subduction without accretion and even extension and normal faulting, may be controlled by relatively small spatial or temporal variations in either  $\mu$  or  $\lambda$ .

J. Geophys. Res., 89, Paper 46080

8150 Plate tectonics: PRESENTIAL RIPPING OF CONTINENTS: A SOURCE OF DISPLACED TERRAINS. Gregory E. Haxel (Department of Geological and Geophysical Sciences, Princeton University, Princeton, New Jersey, 08540), W. Jason Morgan and Hui-Ling Zhou (Department of Geological and Geophysical Sciences, Princeton University, Princeton, New Jersey, 08540). Lithospheric rifting, while prominent in the continents, rarely occurs in oceanic regions. To explain the preferential rifting of continents we compare the total strength of different lithospheres by integrating the effects of lithospheric thickness with depth. Comparisons of total strength indicate that continental lithosphere is weaker than oceanic lithosphere by about a factor of three. A thicker crust can have the total strength of oceanic continental lithosphere. Because the weakest part of the lithosphere is at the surface, the total strength of the lithosphere is controlled by the surface strength. This results in the formation of small continental fragments. The formation of small continental fragments is controlled by the surface strength. This results in the formation of small continental fragments. The formation of small continental fragments is controlled by the surface strength. This results in the formation of small continental fragments.

J. Geophys. Res., 89, Paper 46080

8150 Plate tectonics: PRESENTIAL RIPPING OF CONTINENTS: A SOURCE OF DISPLACED TERRAINS. Gregory E. Haxel (Department of Geological and Geophysical Sciences, Princeton University, Princeton, New Jersey, 08540), W. Jason Morgan and Hui-Ling Zhou (Department of Geological and Geophysical Sciences, Princeton University, Princeton, New Jersey, 08540). Lithospheric rifting, while prominent in the continents, rarely occurs in oceanic regions. To explain the preferential rifting of continents we compare the total strength of different lithospheres by integrating the effects of lithospheric thickness with depth. Comparisons of total strength indicate that continental lithosphere is weaker than oceanic lithosphere by about a factor of three. A thicker crust can have the total strength of oceanic continental lithosphere. Because the weakest part of the lithosphere is at the surface, the total strength of the lithosphere is controlled by the surface strength. This results in the formation of small continental fragments. The formation of small continental fragments is controlled by the surface strength. This results in the formation of small continental fragments.

considerably from the mid-SE structural trends that prevail in the central Red Sea as well as in the faults that border the northern Red Sea and the Gulf of Aden. The tectonic activity that formed the Red Sea rift in the middle-late Miocene was not rejuvenated there during the Pliocene-Quaternary and the new data indicate that the axis of the tectonic extension at

the northern Red Sea jumped to show a northward orientation. It is suggested that an incipient spreading center extends from the Red Sea northwards to the Gulf of Aden and the Dead Sea rift. (Red Sea rift, diapir, spreading center jump).

J. Geophys. Res., 89, Paper 46109

## Chapman Conference on Vertical Crustal Motion: Measurement and Modeling

A Chapman Conference on Vertical Crustal Motion: Measurement and Modeling will be held October 22-26, 1984, in Harpers Ferry, West Virginia.

Convenor: William E. Strange

This conference will bring together scientists who measure vertical crustal motions and those who analyze and model these motions with the primary objective of obtaining close interaction between the two groups. Emphasis will be on vertical crustal motion in North America. Questions to be addressed will be (1) What are the accuracies and error sources associated with each data type? (2) What is the extent of the current data base? (3) How accurately do we know vertical crustal motions in North America? (4) What are realistic expectations of contributions from space systems and other new technologies in the next decade? (5) What is the current status of modeling vertical crustal motions? (6) How important is vertical motion information to understanding and modeling earth dynamics? (7) What are the measurement requirements to support modeling and analysis in terms of temporal and spatial density and accuracy? (8) What are the most critical deficiencies of vertical motion data relative to modeling and analysis?

For housing and registration information contact:  
AGU Meeting Department  
2000 Florida Avenue, N.W.  
Washington, D.C. 20009  
(202) 462-6903For program information contact:  
Dr. W. E. Strange  
NOAA/NOS/CNCS/NGS/NCG11  
601 Executive Boulevard  
Rockville, Maryland 20852  
(301) 443-2520Registration Deadline  
October 1, 1984  
Registration Fee \$75

## News

### Covering for GOES

In the wake of last July's failure of the Geostationary Operational Environmental Satellite (GOES 5) weather satellite over the eastern part of the United States, managers at the National Oceanic and Atmospheric Administration (NOAA) are trying to keep the weather data flowing to satellite users around the country. The geosynchronous satellite, one of two GOES spacecraft stationed over American longitudes, lost its imaging capability on July 29 (Eos, August 21, 1984, p. 483).

Following the failure, the companion GOES-West satellite was shifted from its 135°W station to a more central position over the United States (98°W) so as to cover the eastern part of the country as much as possible. Now, says William Callicott, deputy director of NOAA's Office of Satellite Data Processing and Distribution, the lone remaining U.S. geosynchronous weather satellite will be shifted around with the seasons. At the end of this hurricane season, sometime after November 15, GOES-West (or GOES 6, as it is officially called) will be moved from 98°W to 108°W, where it can better keep watch on winter storms in the northeastern Pacific. Then, in mid-April, after the tornado season, it will be moved back to the central longitude of 98°W.

In the meantime, NOAA is providing weather encoded facsimile (WEFAX) data from non-U.S. geosynchronous satellites (via telephone land lines) to users who normally rely on the GOES data. The European METEOSAT, stationed at 0° longitude, can "see" as far west as 50° and so fills part of the GOES data gap in the Atlantic Ocean. Once the Japanese Geostationary Meteorological Satellite (GMS) is operational in October, it will cover the western Pacific, including Hawaii. By providing WEFAX data from these satellites for areas to the east and west of the North American continent, NOAA hopes to minimize the impact of the GOES failure until a replacement satellite can be launched.

The good news is that the wait for that replacement may be shorter than was previously expected. The spacecraft's builder, Hughes Aircraft Co., hopes to accelerate the production schedule for the next GOES satellite by more than 6 months, according to Callicott. "They're taking all the slack out of the schedule," he told Eos, so that, if everything goes according to "almost a total success schedule," with no unforeseen problems, the satellite would be ready for launch by early autumn of 1985 instead of May 1986.

The bad news is that the wait for that replacement may be shorter than was previously expected. The spacecraft's builder, Hughes Aircraft Co., hopes to accelerate the production schedule for the next GOES satellite by more than 6 months, according to Callicott. "They're taking all the slack out of the schedule," he told Eos, so that, if everything goes according to "almost a total success schedule," with no unforeseen problems, the satellite would be ready for launch by early autumn of 1985 instead of May 1986.

### ESA/NASA Solar Polar Mission Renamed Ulysses

The International Solar Polar Mission (ISPM), a joint project between the European Space Agency (ESA) and NASA, has been renamed "Ulysses." The primary mission of the project, which has not changed with its name, is to investigate the properties of the solar wind, the structure of the sun/wind interface, the heliospheric magnetic field, the interplanetary magnetic field, the solar wind plasma, solar and galactic cosmic rays, and cosmic dust.

Scheduled for launch from the space shuttle in May 1986, the Ulysses mission will make a 14-month journey to Jupiter. That

planet's gravitational effect will deflect the probe into a high-inclination orbit toward the sun. Reaching the sun about 2.5 years later, the satellite will take measurements over the sun's poles. The entire mission will last 5 years.

The mission's new name refers not only to Homer's mythological hero but also to Dante's description in "Inferno" of Ulysses' desire to explore "an uninhabited world behind the sun."

### Glaciology Committee Formed

A committee has been formed to draft a science plan for U.S. involvement in a second deep drilling project on the Greenland Ice Sheet. The first meeting was held in Columbus, Ohio, on July 12, 1984. This committee is addressing two primary issues: (1) potential scientific and logistic involvement in a new multinational program and (2) the advancement of glaciological research in the United States through a science program in Greenland. The committee has received letters from U.S. scientists expressing support and interest in this deep drilling effort.

The members of this committee would be pleased to receive comments or ideas related to possible science programs utilizing the ice core, the borehole, or the logistics facilities at the drill site. The committee members are: Ellen Mosley-Thompson (Chair), Institute of Polar Studies, Ohio State University, Columbus, OH 43210; Anthony J. Gow, U.S. Army CRREL, 73 Lyme Road, Hanover, NH 03755; Michael Herron, Schlumberger-Doll Research, Ridgefield, CT 06877; Kenneth C. Jezek, U.S. Army CRREL, 73 Lyme Road, Hanover, NH 03755; Barclay Kamb, Department of Earth and Planetary Sciences, California Institute of Technology, Pasadena, CA 91109; and Aslam Khalil, Oregon Graduate Center, Oregon State University, Beaverton, OR 97006.

### Geophysicists

Charles Elachi has been named manager of the Earth and Space Sciences Division of the National Aeronautics and Space Administration's Jet Propulsion Laboratory (JPL) in Pasadena, Calif. He succeeds Mustafa Chahine, who assumed the position of JPL chief scientist. Elachi, who joined JPL as a senior scientist in 1971, will continue as principal investigator on the Shuttle Imaging Radar (SIR) series.

Arnold L. Gordon, Professor of Geological Sciences at Columbia University's Lamont-Doherty Geological Observatory, has received the ninth Henry Bryant Bigelow Medal from the Trustees of the Woods Hole Oceanographic Institution in recognition of his 20 years of research in the Southern Ocean and, in particular, for his completion of the Southern Ocean Atlas. In the citation he was described as "the world expert on the physical oceanography of the Southern Ocean."

G. Ross Heath, formerly Dean of the College of Oceanography at Oregon State University, has joined the University of Washington as Dean of the College of Ocean and Fishery Sciences.

Jerome Namias, a research meteorologist at the Scripps Institution of Oceanography, has been awarded the Marine Technology Society's Compass Distinguished Achievement

Award for 1984 "for significant contributions to the art and science of oceanography and marine technology." Associated with Scripps Climate Research Group since 1968, Namias has been involved in long-term weather prediction research and forecasting since the early 1940s. From 1941 to 1964 he served as chief of the Extended Forecast Division of the U.S. Weather Bureau and from 1964 to 1986 as associate director of the National Meteorological Center.

Glenn E. Stout has been named executive director of the International Water Resources Association (IWRA), Urbana, Ill. Stout is currently director of the Water Resources Center at the University of Illinois Urbana-Champaign campus and editor-in-chief of the IWRA publication *Water International*. IWRA's new address is IWRA, University of Illinois, 208 North Romine St., Urbana, IL 61801 (telephone: 217-359-0536).

Alfred M. Fowler has become acting director of the National Science Foundation's Division of Polar Programs.

#### In Memoriam

Ralph M. Berry, 74, died August 22, 1984. A member of the Geodesy Section, he joined AGU in 1959.

Jürgen Henning Jiles, 58, died August 2, 1984. A member of the Tectonophysics Section, he joined AGU in 1973.

### Geophysical Events

This is a summary of *SEAN Bulletin*, 9(8), August 31, 1984, a publication of the Smithsonian Institution's Scientific Event Alert Network. The complete bulletin is available in the microfiche edition of *Eos* as a microfiche supplement or as a paper reprint. For the microfiche, order document E84-009 at \$2.50 (U.S.) from AGU Fulfillment, 2000 Florida Avenue, N.W., Washington, DC 20009. For the paper reprint, order *SEAN Bulletin* (giving volume and issue numbers and issue date) through AGU Separates at the above address; the price is \$3.50 for one copy of each issue number for those who do not have a deposit account, \$2 for those who do; additional copies of each issue number are \$1. Subscriptions to *SEAN Bulletin* are available from AGU Fulfillment at the above address; the price is \$18 for 12 monthly issues mailed to a U.S. address, \$28 if mailed elsewhere, and must be prepaid.

#### Volcanic Events

Kliuchevskoi (Kamchatka): Tephra ejection; lava flows; lahars.  
Mayon (Philippines): Eruption clouds to 15 km; pyroclastic flows; lava flows.  
Soputan (Indonesia): Ash eruption follows local seismicity.  
Api Siau (Indonesia): Tephra ejection, lava flows, lahars; 20,000 evacuated.  
Rabaul (New Britain): Earthquake swarms and slow inflation continue.  
Manam (Bismarck Sea): Volcanic ash explosions; glowing debris avalanches.  
Ulaurun (New Britain): Increased seismicity, then small ash clouds and glow.  
Bagana (Solomon Islands): Lava production continues; SO<sub>2</sub>-rich plume.  
Langila (New Britain): Occasional ash emission; seismicity weak.  
Home Reef (Tonga): Pumice from March-April eruption continues to drift north and west.  
Krafla (Iceland): Eruption from 8.5-km fissure.  
Etna (Italy): Lava production and ash emission continue.  
Lake Monoun (Cameroon): Poisonous gas from lake explosion kills 87.

Pacaya (Guatemala): Summit area lava production continues.

Mount St. Helens (Washington): Intense deformation, then extrusion of new lobe.

Long Valley (California): Seismicity at relatively low level and inflation modest since January 1983 swarm; U.S. Geological Survey says imminent eruption less likely.

Kilauea (Hawaii): Phase 24; high fountains; thin tephra blanket.

Veniaminof (Alaska): Vapor plumes, roaring noise, and felt earthquakes.

Atmospheric effects: 10 years of lidar data from Virginia summarized.

Kliuchevskoi Volcano, Kamchatka Peninsula, U.S.S.R. (56.18°N, 160.78°E). All times are local (= UT + 12 hours).

Eruptive activity which began in March (Eos, p. 425, July 3, 1984) continued through August. During periods of maximum activity, ash was ejected to 5 km and bombs to 1 km above the crater rim (summit elevation 4850 m). Lava flowed to the northwest, northeast, and southwest from the central crater; the largest flow advanced along the northwest valley to about 3 km above sea level and crossed a glacier, forming mud flows. A cinder cone has formed inside the central crater. On August 17 between 0733 and 1027, high-resolution thermal infrared and visual images from polar orbiting weather satellites showed a plume extending about 200 km southeast from the volcano below about 6 km altitude. Soviet volcanologists confirmed these observations, reporting that on August 16-17 a 15-km-wide ash plume extended 200 km from the volcano.

Information Contacts: B. V. Ivanov, Institute of Volcanology, Prib Avenue 9, Petropavlovsk, Kamchatka 683006 U.S.S.R.; Michael Matson, NOAA/NESDIS, Room 510, World Weather Building, Washington, D.C. 20233.  
Soputan Volcano, Sulawesi, Indonesia (1.11°N, 124.73°E). All times are local (= UT + 8 hours).

A 5-hour explosive eruption occurred at Soputan on August 31. This was the first activity since the May 24-26 tephra ejection that deposited more than 10 cm of ash on a 75-km<sup>2</sup> area and forced two airports to close (Eos, p. 425, July 3, 1984).

Volcanological Survey of Indonesia (VSI) seismic instruments recorded a progressive increase in local seismicity beginning August 6. On August 14, a sequence of tremors appeared between 0400 and 0800, with amplitude increasing to 25 mm (at 2000 magnification). VSI issued a warning to civil authorities and an alert was put into effect on August 14. Seismicity continued August 15-25 with an irregular number of A- and B-type events, averaging 1-2 per day. From August 25 until the time of the eruption, seismicity totally stopped, increasing suspicion among VSI scientists that an eruption was possible.

The eruption started at 0709 on August 31 and lasted until about noon. An ash column rose to about 6 km and moved northeastward. Authorities and area residents were well prepared, and neither casualties nor an evacuation were reported. Press sources reported that the ash cloud could be seen from Manado, the provincial capital 80 km to the north. The ash cloud covered a large area and disrupted traffic on the Trans-Sulawesi highway.

On August 31 at 1457, a visible band image from the NOAA 7 polar orbiting satellite showed a plume extending about 450 km westward from the volcano. The plume was quite dense and about 120 km high.

Information Contacts: Adjat Sudradjat, Director, Volcanological Survey of Indonesia, Diponegoro 57, Bandung, Indonesia; Michael Matson, NOAA/NESDIS, Room 510, World Weather Building, Washington, D.C. 20233; United Press International.

Krafla Caldera, Mývatn Area, Iceland (63.73°N, 16.68°W). The following is a report from Karl Grönvald and Páll Einarsson.


"After a quiet interval of 2 years and 9 months, an eruption broke out at Krafla on September 4, 1984. The eruption in November 1981 and associated deflation of magma reservoirs below Leirhnjúkur were followed by inflation, reaching previous levels in early 1982. Since then slow and intermittent inflation has continued, accompanied by earthquakes in the reservoir roof.

"Rapid deflation over the magma reservoir, followed by volcanic tremor began September 4, at about 0200, but the eruption broke out at 2349. The beginning of the eruption was observed from the air by alerted scientists and a television reporter. The first fissure segment opened about 6 km N of Leirhnjúkur, followed within a minute by another about 3 km to the S. The fissures quickly joined and in 1 hour reached their full length of 8.5 km, extending from Leirhnjúkur to the N. During the first hours lava was erupted along the whole fissure, advancing on broad fronts.

"Already in the early hours of the morning, the activity had decreased and lava production on various sections of the fissures had faded out. By midday on September 5, inflation had resumed. By September 6,

News (cont. on p. 733)





The focal point for physical, chemical, geological, and biological oceanographers.

Editor: David A. Brooks, Department of Oceanography, Texas A&M University, College Station, TX 77843 (telephone: 409-845-5527).

## Drifting Derelict Trajectories in the North Atlantic

Philip L. Richardson

### Introduction

In December 1883 the U.S. Navy Hydrographic Office, a branch of the Bureau of Navigation of the Navy Department, began to publish monthly Pilot Charts. Earlier, oceanographer M. F. Maury had produced some summary survey charts showing ocean currents, winds, sailing routes, and the locations of whales. The new charts were unique in that they showed updated positions of derelict vessels and other drifting debris. From this series of positions of identified derelicts the first ocean trajectories were obtained. Much of this information has been forgotten during the last 100 years, and good collections of the Pilot Charts are rare. (The only complete collection that I could find is held by the Defense Mapping Agency.) This article is a recompilation and description of these early trajectories and a reminder of the usefulness of the Pilot Charts. It also provides a glimpse of a little known part of maritime history, the last days of wooden sailing vessels.

### The Pilot Chart

The new Pilot Charts were prepared to supply a reliable plotting sheet and a graphic presentation of recent as well as general summary information for mariners [Hayden, 1888]. The charts were issued free to navigators in return for their reporting recent navigational and weather information. The success of the Pilot Chart was due to the large number of observers that contributed information for each month's chart and its rapid distribution to ships. In the late 1800's there were nearly 3000 voluntary observers, mariners who crossed the North Atlantic and who patrolled its waters. Reports of marine meteorology and dangers to navigation were collected from these observers at 11 branch offices and forwarded to headquarters in Washington where the latest positions of wrecks and derelicts were plotted on a large blackboard. The Pilot Charts incorporated these data and were published and distributed at the beginning of each month. In November 1893 the office received no less than 400 reports daily from vessels in the North Atlantic alone. In New York during 1886-1887, 6,739 vessels were visited, 3,601 reports were forwarded to Washington, nautical information was furnished to 83,345 masters of vessels and others, and 10,397 Pilot Charts were distributed [Hayden, 1888].

### Derelicts

A derelict is a vessel abandoned at sea. Derelicts that survived more than a few days at sea were usually wooden sailing vessels, and the longest surviving of these were often lumber schooners. From the point of view of a ship captain, a derelict vessel is a formidable obstruction to navigation. A collision with a derelict at night or in fog could damage or sink a ship. In our age of metal ships it is not generally recognized how many derelicts there were nor how long they remained afloat. The Atlantic was literally strewn with numerous "Mary Celestes" in various stages of disintegration.

The number of reported derelict sightings reached a maximum toward the end of the 19th century [Hydrographic Office, 1894]. During 1893, a year of particularly numerous derelicts, there were 782 reports of 418 different derelicts. One hundred six of these derelicts were identified by name. All but two

or three of these derelicts were wooden. Over a 7 year period, 1887-1893, a total of 1,628 derelicts were sighted, an average of 232 annually, 19 per month. This suggests that at any one time at least 19 derelicts remained afloat in the North Atlantic. The average length of time a derelict remained afloat is estimated to be 30 days. This time is based on assuming a derelict remained afloat 1 day after its last reported sighting or 5 days for a single sighting. During 1887-1893 there were 1,944 reports of the 482 identified derelicts giving an average of four sightings per derelict. The greatest number of derelicts were first reported in September-November and were caused by severe storms. Most were located in the Gulf Stream off the U.S. coast. The numbers of sightings gradually decreased eastward along the transatlantic steamer routes. Many of the large number of derelicts observed during the fall of 1893 were caused by a series of three hurricanes which occurred in August of that year.

### Derelict Trajectories

Numerous derelicts remained afloat over half a year and were reported often enough to give long and interesting ocean trajectories. A listing of derelicts that floated longer than 200 days is given by Richardson [1984]. Six of the derelicts drifted longer than a year: (1) schooner *Fannie E. Walton*, 1100 days; (2) schooner *Wyer C. Sargent*, 615 days; (3) bark *Trensch*, 551 days; (4) bark *Vincenzo Perrotta*, 536 days; (5) schooner *Ethel M. Davis*, 370 days; and (6) schooner *James B. Drury*, 367 days.

The trajectories of three long-lasting and far-drifting derelicts are shown in Figure 1. One of the best known of these was the three-masted lumber schooner *W. L. White*, belonging to A. F. Ames of Rockland, Maine. She was abandoned off Delaware Bay during the great blizzard of March 15, 1888. A telegram dated Stornoway, Hebrides Islands, Scotland, January 25, 1889, marked the termination of the *White*'s 310 day transatlantic drift. She ended strand on Heaskey Island in the Hebrides.

The *White* began her drift southward under the influence of the inshore current and northwest gale, with masts and portions of her sails standing. Upon reaching the Gulf Stream she turned and followed a east-northeast course at an average speed of about 32 miles per day. From May to November 1888 she looped and zigzagged east of Newfoundland directly within a major shipping lane. During these 6 months she was reported by 36 vessels, three of which sighted her in a single day. In her cruise of 10 months and 10 days, she traversed a distance of 5,900 miles and was reported 45 times.

Although the detailed paths of the derelicts are very different from each other, there are some similarities which might be described as patterns. Eight of the longest drifting derelicts moved eastward in the Gulf Stream until they reached 50°W where their paths diverged. Three derelicts continued eastward and crossed the Atlantic in an average time of 10 months. The *White* took 310 days, the *Twenty-one Friends* took 255 days, and the *Hunt* took 347 days. Six derelicts drifted southward from the Gulf Stream near 40°W. The *Drury* and the *Hill* both made light turns and drifted westward just south of the Gulf Stream. The *Walton* made a complete circuit of the gyre during its 3-year drift. This derelict drifted south to 25°N, westward to the Bahamas, and then northward into the Gulf Stream again, crossing its earlier path. The trajectory of the *Trensch*, which was 1.5 years long, is similar to part of the *Walton*'s. Two derelicts drifted erratically but in a general southward direction through the Sargasso Sea and grounded on the Bahama Islands.

Most derelicts looped as they drifted. The *Sargent* and *Walton* made large, 500 km loops with a characteristic period of 10 months near 30°N, 40°W. Several other derelicts made frequent smaller scale loops: the *Perrotta* and *Francis* in the Sargasso Sea and the *White* east of Newfoundland.

An example of variability of ocean surface currents is given by the drift of the bow and stern of the *Fred B. Taylor*. On June 22, 1892, the *Traue* collided with the *Taylor* and the latter was cut in two (from Pilot Chart, September 1892). The forward and after parts separated and drifted in entirely different directions (Figure 2). The bow went 340 miles during 93 days and was reported 47 times. The stern went 350 miles during 47 days and was reported 20 times. The different directions could have been partly caused by the different areas of bow and stern presented to the wind and current.

### Superimposed Trajectories

A summary diagram was prepared that shows 200 derelict trajectories reported in the Pilot Charts from 1883 to 1902 (see cover). Earlier but less complete charts showing trajectories of derelicts have been given in the supplements of February, 1889 and 1893 to

the Pilot Charts, by the Hydrographic Office [1894], and by *Hautreux* [1897]. Derelict vessels which first appeared near the U.S. Coast south of Long Island and north of Cape Hatteras usually drifted in a southward direction following the inshore current until they reached Hatteras, where they entered the Gulf Stream and drifted eastward.

In general, derelicts entered the Gulf Stream north of 30°N and moved eastward in the Stream. When they reached the area south of the Grand Banks, near 40°N, 50°W, they split into two bands of trajectories. The first band reaches northeastward and then eastward, passing north of the Azores between 40° and 50°N. The second band extends southeastward and then westward near 25°N. Six derelicts moved southward between the Azores Islands and Spain and Portugal. The general pattern indicated by the collected trajectories is of a large clockwise gyre split into two branches, one branch located north of the Azores, the other southwest of the Azores. The splitting of the Gulf Stream near the Grand Banks has been confirmed by more recent measurements [Mann, 1967; Clarke et al., 1980], but it is still controversial [Worthington, 1976].

Superimposed on the large-scale, long-term general circulation pattern can be seen considerable current variability. The derelicts do not often smoothly follow the large-scale gyre; instead they drift in convoluted trajectories that often cross each other. The convoluted paths give an early Lagrangian measure of mesoscale eddies and longer period current fluctuations. We now know that the ocean is populated by energetic eddies that are usually much stronger than the mean currents [Schmitz et al., 1983; Robinson, 1983]. Recently, the importance of these eddies to the general circulation has been recognized, and they have been studied intensively. Because of these eddies, the mean circulation

becomes recognizable only by averaging a great quantity of observations in space and time, as can be done by eye on the cover. In the Gulf Stream, the North Atlantic Current, and the North Equatorial Current, one clearly sees the general drift in spite of the eddies. In the Sargasso Sea the trajectories are dominated by mesoscale eddy motion.

One should be cautious about interpreting all the motion indicated by trajectories as being due to water movement. Derelict ships varied in size and weight and in state of damage when abandoned. Some were totally dismasted and filled to the gunwales with water. Along with the 30% of the sightings which were of vessels that had turned bottom up, these probably provided a good indication of the speed and trajectory of near surface water. Derelicts with masts standing and those riding high in the water would no doubt be significantly influenced by the winds blowing directly on the mast and exposed hull.

There are the additional problems of position errors of the reporting ship, misidentification of derelicts, and infrequent sightings. It is difficult to evaluate with the available information how accurate the details of trajectories really are. The average number of days between sightings is about 20, which is sufficiently small that we can see some aspects of mesoscale eddies but not small enough except in a few occasions to resolve individual loops. Despite the lack of frequent and precise positions, the collection of the trajectories represents a realistic view of the general Lagrangian circulation and of current variability.

Within the last decade, freely drifting buoys remotely tracked by satellite have begun to be used in large numbers to measure velocities and trajectories of near-surface currents. The newer measurements have the advantage over earlier derelict trajectories of several fixes per day and a higher positional accuracy. From a collection of these measure-




Fig. 1. Trajectories of (1) the schooner *W. L. White* from March 15, 1888, to January 25, 1889, a drift of 5,190 miles and 310 days; (2) the schooner *Wier C. Sargent* from March 3, 1891, to December 6, 1892, a drift of 5,500 miles and 615 days; and (3) the bark *Vincenzo Perrotta* from September 17, 1887, to April 4, 1889, a drift of 2,960 miles and 536 days.




Fig. 2. Drift of the bow and stern of the *Fred B. Taylor*.

ments we have been able to obtain a more quantitative picture of aspects of the general circulation and of the geography of ocean variability. A summary figure of the buoy trajectories [Richardson, 1983] shows general patterns very similar to those of the derelict trajectories. If these buoys continue to be deployed in the North Atlantic we might expect that the numbers of their trajectories may eventually surpass the numbers of drifting derelict trajectories.

Derelict sightings and trajectories have been viewed here as giving interesting information about ocean currents. We should not forget that this information came with a tragic loss of life and shipping. Toward the end of the 19th century, 12,000 lives and 2,200 vessels were lost at sea each year worldwide (supplement to February 1893 Pilot Chart). Each severe storm that was encountered at sea left new derelicts in its path and added new names to the long list of vessels and men who left port but were never heard of again. The plots of derelict sightings are a sad reminder of lost vessels, suffering, and death.

### Summary

Pilot Charts, published monthly during the last two decades of the 19th century, reveal a rare, interesting, and tragic glimpse of maritime history in detailing observations of drifting derelict ships. The large collection of derelict sightings was made possible by the excellent and fast reporting system established by the Navy Hydrographic Office. Numerous voluntary observers reported dangers to navigation which were quickly incorporated into the next month's chart. By 1900, however, wooden sailing ships, which comprised most of the derelicts, had been superseded by steamers, and derelicts became infrequent. Drifting derelicts gave some first examples

of ocean trajectories. They showed the general pattern of circulation in the North Atlantic Ocean including the bifurcation of the Gulf Stream near the Grand Banks of Newfoundland. The trajectories gave an early indication of current variability. Coupled with set and drift measurements from ships underway, these derelict trajectories provided much of the early knowledge of near-surface ocean currents.

### Acknowledgments

A. Green kindly made available the Defense Mapping Agency's collection of Pilot Charts, 1883-1902. Funds were provided by the National Science Foundation under grant OCE81-09145. This is an abbreviated version of a paper that discusses the same information [Richardson, 1984].

### References

- Clarke, R. A., H. W. Hill, R. F. Reiniiger, and B. A. Warren, Current system south and east of the Grand Banks of Newfoundland, *J. Phys. Oceanogr.*, 10, 25-65, 1980.
- Hautreux, M., *Atlantique Nord-Courants de Surface*, Congrès National des Sociétés Françaises de géographie, Société de Géographie Commerciale de St. Nazaire, Imprimerie Fronteau, St. Nazaire, 1897.
- Hayden, E., The Pilot Chart of the North Atlantic Ocean, *J. Franklin Inst.*, 125, 265-278, 447-462, 1888.
- Hydrographic Office, *Wrecks and Derelicts in the North Atlantic Ocean, 1887 to 1893*, Inclusive, Government Printing Office, Washington, D.C., 1894.
- Mann, C. R., The termination of the Gulf Stream and the beginning of the North Atlantic Current, *Deep Sea Res.*, 14, 337-359, 1967.
- Richardson, P. L., Eddy kinetic energy in the North Atlantic from surface drifters, *J. Geophys. Res.*, 88, 4355-4367, 1983.
- Richardson, P. L., Drifting derelicts in the North Atlantic 1883-1902, *Prog. Oceanogr.*, in press, 1984.
- Robinson, A. R. (Ed.), *Eddies in Marine Science*, Springer-Verlag, New York, 1983.
- Schmitz, W. J., Jr., W. R. Holland, and J. F. Price, Mid-latitude mesoscale variability, *Rev. Geophys. Space Phys.*, 21, 1109-1119, 1983.
- Worthington, L. V., On the North Atlantic Circulation, *The Johns Hopkins Oceanogr. Stud.*, vol. 6, Baltimore, Md., 1976.

Philip L. Richardson is with the Woods Hole Oceanographic Institution, Woods Hole, Mass.

## Mathematical Models for Zooplankton Swarms: Their Formation and Maintenance

Akira Okubo and James J. Anderson

### Introduction

Many aquatic invertebrates are known to form swarms and schools. Thus, zooplankton are usually distributed unevenly both in vertical and in horizontal directions. There are a great number of studies on zooplankton swarms ranging from simple records or observations to more extensive functional and behavioral investigations. Yet, no attempt has been made on mathematically modeling these phenomena, chiefly because of the lack of detailed data on individual movements in swarms.

Swarms are groups of individuals engaging in more or less cohesive movements without parallel orientation. The presence or absence of parallel orientation distinguishes schools from swarms. Individual organisms in a swarm apparently exhibit irregular movements which might be regarded as random. However, random motion alone makes a group of organisms spread out to occupy a larger space as time progresses (i.e., diffusion). Therefore, an adequate model for zooplankton swarming must assume certain regularities in the motion of individuals superimposed upon its randomness. This deterministic part arises primarily from behavioral interactions between swarming individuals and possesses the nature of an attractive force. A swarm is maintained by the balance between the deterministic and stochastic forces. Modeling for swarm formation involves random search for prey or conspecific individuals followed by accelerated aggregation as a result of mutual communication or biological cue.

Does your library subscribe to the Water Resources Monograph Series?

In this report we will present both kinematic and dynamical models for the maintenance and formation of swarms.

### Kinematic Interpretation of Difference Between Swarming and Diffusion

We consider a simple one-dimensional case (x axis). Swarming implies randomness in motion. Thus both velocity  $v(t)$  and displacement  $x(t)$  of an individual in a swarm are random variables.

As a statistical measure determining the swarm dimension, the variance of the displacement is calculated by averaging the square of  $x$  over the ensemble (e.g., over a sufficiently large number of individuals in a swarm). Without loss of generality we assume that the individual organisms to start at  $x = 0$  and the individual movement to be symmetrical with respect to the origin so that the swarm centroid remains at the origin. Thus, the variance of  $x$  is obtained by

$$\overline{x^2}(t) = \int_0^t \langle v(t')v(t') \rangle dt' = 2 \int_0^t \int_0^{t'} \tau R_L(\tau) d\tau dt = 2 \int_0^t \tau R_L(\tau) dt$$

where

$$R_L(\tau) = \langle v(t')v(t'+\tau) \rangle / v^2$$

is the Lagrangian velocity autocorrelation coefficient. We have assumed that the random velocity field is stationary so that the correlation coefficient depends only on the time lag  $\tau$ .

Equation (1) provides us with kinematic distinction between diffusion and swarming in terms of the velocity autocorrelation. By diffusion we mean that the individual velocity loses its statistical dependence on past velocities as the dispersion continues. In other words,  $R_L(\tau)$  approaches zero at large time lags such that the limit integral in (1) converges as  $t \rightarrow \infty$ . We then find from (1) that as  $t \rightarrow \infty$ ,

$$\overline{x^2}(t) \rightarrow 2Dt$$

where

$$D = \int_0^\infty \tau R_L(\tau) d\tau$$

If, on the other hand, (1) is to mathematically describe swarming,  $\overline{x^2}$  must approach a constant value for steady state swarm maintenance. This is satisfied only if  $R_L(\tau)$  oscillates about zero value in such a manner that  $\tau R_L(\tau)$  approaches zero asymptotically, and, consequently, the second integral of (1) approaches a negative constant value. In physical terms, the individual motions appear to resemble a random pendulum-like motion about  $x = 0$ . Experimental confirmation on this model would be to determine the velocity autocorrelation coefficient of individual animals in a swarm.

### Dynamical Model for the Maintenance of Swarm

We now consider a dynamical model for swarm maintenance. Newton's equation of motion will be applied to swarming animal motion. We assume that (1) the frictional force is proportional to the velocity of organism (in fact, Reynolds numbers associated with zooplankters range from 0.1 to 500 [Zaret, 1980] so that the Stokes' law of drag may not be applicable to some large and fast moving zooplankton); (2) the nonrandom force is attractive by nature toward the center of swarm and dependent on the distance from the center; and (3) the random force is a type of white noise.

Then the equation of motion is given by

$$m \frac{d^2x}{dt^2} = -k \frac{dx}{dt} - \omega^2 x - \phi(x) + A(t)$$

where  $k$  is the frictional coefficient,  $\omega$  is the frequency of harmonic component of the attractive force,  $\phi(x)$  is the acceleration due to anharmonic component of the attractive force, and  $A(t)$  is the random acceleration of a white noise nature;

$$\overline{A} = 0 \quad \langle A(t)A(t+\tau) \rangle = 2B\delta(\tau) \quad (6)$$

where  $B$  is the intensity of the variance of  $A(t)$ .

An approximate analytical solution of (5) may be obtained by the method of equivalent linearization [Bulawa et al., 1982]. To this end, (5) is replaced by a linear system with an equivalent linear frequency  $\omega_e$  such that

$$\omega_e^2 = \omega^2 + \langle \phi(x) \rangle / x^2 \quad (7)$$

where the angle brackets denote a time average over one cycle of oscillation of  $x = a \sin \omega t$ .

Solving this linearized version of (5) for  $x$ ,

we calculate the velocity-autocorrelation coefficient to be

$$R_L(\tau) = e^{-k\tau/2} (\cos \omega_e \tau - k/2\omega_e \sin \omega_e \tau) \quad (8)$$

where

$$\omega_e^2 = \omega^2 - k^2/4 \quad (9)$$

Also the variance is found to be

$$\overline{x^2} = B/k\omega_e^2 \quad (10)$$

as  $t \rightarrow \infty$ . This dynamical model enables us to obtain the variance of displacements of swarming animal in terms of parameters that characterize the motion of individuals. The square root of (10) scales the spatial extent of swarm. With certain modifications on the stochastic forcing function  $A(t)$  the dynamical model equation (5) or its equivalent linearization is also applicable to schooling of fish [Okubo, 1980].

In view of the assumed nature of the stochastic force the random process  $(x, v)$  is Markovian, and the probability density function (p.d.f.)  $p_2(x, v, t)$  obeys a Fokker-Planck equation. We thus obtain from (5) and (6)

$$\frac{\partial p_2}{\partial t} = -v \frac{\partial p_2}{\partial x} - \frac{\partial}{\partial v} \left\{ (-kv - \omega^2 x - \phi(x)) p_2 \right\} + B \frac{\partial^2 p_2}{\partial v^2} \quad (11)$$

Since  $p_2$  is an even function of  $x$  and the non-random attractive force is an odd function of  $x$ , integral of (11) over  $x$  leads to

$$\frac{\partial p_1}{\partial t} = k \frac{\partial}{\partial v} (v p_1) + B \frac{\partial^2 p_1}{\partial v^2} \quad (12)$$

where

$$p_1(v, t) = \int_{-\infty}^{\infty} p_2(x, v, t) dx \quad \text{p.d.f. of velocity} \quad (13)$$

For steady state  $p_1^*$

$$p_1^*(v) = (k/2\pi B)^{1/2} e^{-k^2 v^2 / 2B} \quad (14)$$

which is the Maxwellian velocity distribution. The p.d.f. of displacement  $x$  may be obtained by integrating  $p_2(x, v, t)$  over  $v$

$$p(x, t) = \int_{-\infty}^{\infty} p_2(x, v, t) dv \quad (15)$$

Alternatively, if we use the Smoluchowski-Kramers approximation or the method of adiabatic elimination [Gardiner, 1983] in (5), we obtain

$$\frac{\partial p_1}{\partial t} = \frac{\partial}{\partial x} \left( \frac{\omega^2 x + \phi(x)}{k} p_1 \right) + B \frac{\partial^2 p_1}{\partial x^2} \quad (16)$$

The steady state p.d.f.  $p_1^*$  gives the number density distribution of swarming organisms

$$p_1^*(x) = p_0 \exp \left\{ -\frac{\omega^2 x^2}{2B} - \int_0^x \frac{\phi(x)}{B} dx \right\} \quad (17)$$

where  $p_0$  is the density at the swarm center. The non-Gaussian density distribution arises from the anharmonic attractive force. Since  $\phi(x) > 0$ , the density distribution tends to be platykurtic. It is conceivable that the anharmonic attractive force is a manifestation of density-dependent advective velocity toward the swarm center. It is a built-in mechanism to maintain a sharp boundary of concentration despite a general tendency to spread by the random component of motion. The net effect also produces a more or less uniform density of organisms within a swarm which is characteristic of a platykurtic distribution.

### Comparison With Data

Verification of our mathematical models for swarming requires comparisons with appropriate data; to our regret no such data really exist in the marine field. In desperation we are forced to borrow data from "terrestrial" (i.e., insect) swarming.

High-speed filming on swarming of midges, *Anopheles tritaeniorhynchus* Kim, provides us with basic kinematic data for evaluating swarm maintenance characteristics such as velocity autocorrelation coefficient, velocity frequency distribution, acceleration field, swarm dimensions, and insect number density distribution [Okubo and Chiang, 1974; Okubo et al., 1977].

Figure 1 shows the velocity- and acceleration-vector field of individual midges in an instance of swarming. The midges of size 1-2 mm moves with high acceleration, occasionally twice or more the acceleration of gravity ( $g$ ). The motion inside the swarm is more or less random in both velocity and acceleration, but the midge is subject to an inward force (or rather acceleration) which is felt strongly at the swarm edge. As we model in (5), the presence of this inward-oriented force plays a crucial role in maintaining the

Oceanography (cont. on p. 732)






Fig. 1. Velocity and acceleration field of individual animals in an instance of swarming.




Fig. 2. Mean acceleration of swarming animals versus normalized distance from the swarm center.




Fig. 3. Velocity autocorrelation coefficient of swarming animals versus time lag.




Fig. 4. Speed frequency distribution of swarming animals.

swarm against a general tendency of spread by the randomness in the motion. The acceleration field in swarming can be seen more clearly in Figure 2, where we plot the mean acceleration of swarming animals against normalized distance from the swarm center.

Figure 3 shows the velocity autocorrelation coefficient against time lag. The theoretical curve is based on (8) with  $k = 6.8/s$  and  $\omega_p = 31.6/s$ . The observed behavior of the autocorrelation coefficient is consistent not only with the kinematic interpretation of swarming but also with the dynamical model.

Figure 4 shows the frequency distribution of insect speed in a swarm, which can be well approximated by the Maxwell speed distribution. The theoretical curve is based on (14) with  $k = 6.8/s$ ,  $B = 1.22 \times 10^4 \text{ cm}^2/s^2$ , the value of  $B$  is estimated from the observed mean squared velocities of individuals.

The number density distribution of swarming individuals is in fact platykurtic as our mathematical model, equation (17) predicts. For six swarms of *Anastrepha* the mean kurtosis is 2.60, and the difference from normality (kurtosis = 3) is statistically significant. As explained earlier, this deviation from normality arises from anharmonic attractive forces that correspond to density-dependent advective velocity toward the swarm center. Okubo and Chiang (1974) obtained empirically with *Anastrepha* swarms that the advective velocity scaled by diffusivity is proportional to the minus  $1/4$  power of the number density.

## Dynamical Model for the Formation of Swarm

Modeling swarm formation is much more difficult than that of swarms maintenance simply because we deal essentially with a transient behavior. The dynamical model of swarm formation assumes two processes (i.e., initial random encounter of conspecific individuals and accelerated aggregation followed).

The initial phase of swarm formation may be modeled in a way similar to Anderson (1981). Namely, the change with time of the number of individuals  $N(t)$  in a swarm is determined as the difference between the rate that zooplankton enter the swarm, which is independent of  $N$  without communication, and the rate that zooplankton exit from the swarm, which is proportional to  $N$ . We thus have

$$\frac{dN}{dt} = a - bN \quad (18)$$

where  $a$  and  $b$  are constants. Possible stochasticity in  $a$  and  $b$  may be incorporated (Anderson, 1981). The solution of (18) subject to  $N = 1$  (one individual) at  $t = 0$  describes an asymptotic approach to a stable equilibrium  $N_e = a/b$  with a rate of  $b^{-1}$ .

This equilibrium state would not be realized, however, if mutual communication exists as swarming progresses. Some possible mechanisms by which zooplankton might communicate are light, pressure changes due to sound generation, and chemosensory by means of pheromones. Aggregation behavior in response to pheromones has been well known in insects, and chemical communication has been observed in copepods and planktonic shrimp.

If chemosensory is assumed as the mechanism of zooplankton communication, we can

construct a simple model for the accelerated aggregation process that leads to a massive swarm or "superswarm."

First consider diffusion of a chemical from a spherical source of radius  $R$ . For steady state the concentration of chemical  $C$  is given by

$$C(r) = \frac{Q}{4\pi Dkr} \quad r \geq R \quad (19)$$

where  $Q$  is the rate of release of chemical,  $D$  is diffusivity, and  $r$  is the distance from the center of the sphere.

We now regard the source as a swarm, in which  $N(t)$  is the total number of individuals,  $p$  is the mean volume occupied by one individual, and  $q$  is the emission rate of chemical by one individual;

$$C(r, t) = \frac{qN(t)}{4\pi Dkr} \quad (20)$$

Let  $C^*$  be the threshold concentration for chemosensory response. Then the effective sphere of chemical communication around the swarm has the volume of

$$V = \frac{4}{3}\pi r^{*3} = \frac{4}{3}\pi R^3 \left( \frac{qN}{4\pi DkC^*} \right)^3 = \frac{4}{3}\pi \left( \frac{qN}{4\pi DkC^*} \right)^3 N^3 = pN^4 \quad (21)$$

Any individual who happens to be within the sphere of influence will be attracted to the swarm by chemotaxis. Let  $v$  be the velocity of chemotaxis and  $S_0$  be the background density of zooplankton. Then the change with time of the number of individuals in the swarm is given by

$$\frac{dN}{dt} = v_0 V = v_0 \mu (N^4 - \lambda N) \quad (22)$$

where

$$\mu = \frac{4\pi}{3} \left( \frac{qS_0}{4\pi DkC^*} \right)^3$$

and

$$\lambda = \frac{3}{4\pi} \left( \frac{4\pi DkC^*}{q} \right)^3$$

The solution of (22) subject to  $N = N_0$  at  $t = t_0$  is obtained by

$$N(t) = \left[ \frac{\lambda}{1 - (1 - \lambda N_0^{-3}) \exp(2\lambda \mu S_0 (t - t_0))} \right]^{1/3} \quad (23)$$

This solution describes the avalanche of aggregation in such a manner that the swarm reaches an infinite size within a finite time span given by

$$t - t_0 = (2\lambda \mu S_0)^{-1} \ln(1 - \lambda N_0^{-3})^{-1} \quad (24)$$

which for small  $\lambda N_0^{-3}$  approximates

$$t - t_0 = (2\lambda \mu S_0)^{-1} \quad (25)$$

The knowledge of the pertinent parameters would enable us to estimate this critical time. Again, no such data exist in marine swarming. The presented model may also be applicable to acoustic communications with minor modifications.

## Discussion

We have outlined mathematical models for the formation and maintenance of zooplankton swarms. The theories extend beyond the ability of observation in many instances, and this may be criticized since some consider that theory without observations is a scientific crime. In fact we believe it is a crime not to. In justification of this view we reference the synergistic relationship between theory and observation in particle physics. In many ways, zooplankton are not unlike particles but with very elaborate laws of governing their behavior.

Our approach at this time is to consider the very basic features of swarming and to compare our mathematical models with available data on insect swarming. To proceed into marine swarming, kinematic data on zooplankton swarming are urgently needed for evaluation of the mathematical models. For example, information on the basic swarm properties: the density of swarms in a habitat and the density of zooplankton in a swarm will enable us to estimate at least the ratio of the diffusivity and the attractive forces in terms of the number of density (Okubo and Chiang, 1974).

In perspective, this paper is written by two persons who eagerly undertake to build models of zooplankton swarming in the dim hope that they will find a scent of a "working hypothesis" on swarming that can be tracked by biologists. It remains to be determined if the creature whose track we follow will be categorized as "BIOSENSIS" or "BIONONSENSIS."

## References

- Anderson, J. J., A stochastic model for the size of fish schools, *Fish. Bull.*, 79, 315-323, 1981.
- Bulsara, A. K., Lindenberg, K. E., Shuler, Application of linearization methods to driven nonlinear systems, in *Instabilities, Bifurcations, and Fluctuations in Chemical Systems*, edited by L. E. Reichl and W. C. Schieve, pp. 400-410, University of Texas Press, Austin, Texas, 1982.
- Gardiner, C. W., *Handbook of Stochastic Models*, Springer-Verlag, Berlin, 1985.
- Okubo, A., *Diffusion and Ecological Problems*, Springer-Verlag, Berlin, 1980.
- Okubo, A., and H. C. Chiang, An analysis of the kinematics of swarming of *Anastrepha pritchardi* Kim (Diptera: Cecidomyiidae), *Res. Popul. Ecol.*, 16, 1-42, 1974.
- Okubo, A., H. C. Chiang, and C. C. Ebbesmeyer, Acceleration field of individual midges, *Anastrepha pritchardi* (Diptera: Cecidomyiidae), within a swarm, *Can. Entomol.*, 109, 149-156, 1977.
- Zaret, R. E., The animal and its viscous environment, in *Evolution and Ecology of Zooplankton Communities*, edited by W. C. Kerfoot, pp. 3-9, University Press of New England, Hanover, N.H., 1980.

Akira Okubo is with the Marine Sciences Research Center, State University of New York, Stony Brook, NY 11794.

James J. Anderson is with the Fisheries Research Institute, University of Washington, Seattle, WA 98195.

## Information Report

### Blowatt: A Study of Bioluminescence and Optical Variability in the Sea

On entering water, light is both scattered and absorbed. The sum of these is attenuation. In the open ocean, the agent of most scattering and absorption is plankton, or plankton-derived products. The controls, therefore, governing the variability in light absorption and scattering are no less than the controls governing the distribution, abundance, and growth of plankton populations. Apart from being attenuated, light is also generated ubiquitously in the upper layers of the sea through mechanisms of bioluminescence. The overall and long-term goal of Blowatt is to establish causal links operating between the variability in light attenuation and light production in the ocean. The issues addressed range from behavioral relationships among macrozooplankton and micronekton, to the dynamics of absorbing and scattering populations, to the physical dynamics of the upper layers. A conceptual model is shown in Figure 1.

The beginnings of Blowatt can be traced to an advisory meeting convened by the Office of Naval Research (ONR), in December 1982 at Berkeley, Calif. The 30-odd meeting attendees explored various research and instrumentation development avenues within the framework of bioluminescence and optical variability in the ocean. Preliminary recommendations from the group covered immediate tasks, overall design of the program, possible field sites, and the sort of modeling effort which should accompany and guide the observations (Blizard et al., 1982). During

1983, the program continued to evolve in concept, drawing much scientific support and insight from a predecessor ONR program, the Optical Dynamics Experiment (ODEX). Funding approval from ONR for Blowatt came in early 1984, at which time a steering committee was selected. (The Blowatt Steering Committee members are James F. Case (UCSB), Mary Jane Perry (UW), Raymond C. Smith (UCSB), and Elijah Swift (URI).) The program is expected to have a duration of 5 years, with field years in 1985 and 1987. We now turn to some of the specific issues Blowatt will address in the coming field experiments.

### Optical Variability in the Ocean

Biological and optical oceanographers traditionally deal with plankton populations in terms of such macroscopic, but easily measured, variables as chlorophyll *a*, particle counts and beam transmittance. In many kinds of biological oceanographic studies, such as the influence of physical processes on phytoplankton, these are the variables of choice since the sampling design necessitates near continuous data acquisition. However, studies of the dynamics of plankton communities require much more elaboration of organism type and function, and trophic processes, but with a compromise of sampling coverage. An understanding of the optical variability of the ocean will require both kinds of research: the physical forcings upon macroscopic variables as well as the detailed character of the biological distributions and dynamics that these forcings permit.

For example, at one level, we need to understand how physical processes in the upper ocean influence vertical distributions of chlorophyll *a* and particulate matter, and thereby the behavior of inherent and apparent optical properties. (These optical properties refer, respectively, to those independent of and dependent on, light source distribution.) A variety of physical processes contribute to the mass, momentum, and energy budgets in the upper ocean, however, a few phenomena can be identified as major sources of the variance spectra of physical and biological data fields. According to Klein (1980) and Dickey and Simpson (1983), wind and surface current interactions caused by inertial resonance phenomena may explain rapid mixed-layer deepening, which, if true, means that nutrient fluxes may vary substantially on time scales of hours (Klein and Coste, 1984).

Beyond this, developing predictive relationships between biological particles and optical properties in the ocean requires much more exhaustive specification of the nature of absorbers and scatterers. From an optical perspective, biogenic particles are ensembles of sizes, shapes, pigments, and refractive indices. We do not understand enough about the nature of the particles themselves and their distributions (let alone their dynamics) to enable an understanding of their influences to variability in absorbance and scattering.

Although the various types of phytoplankton (diatoms, dinoflagellates, cyanobacteria, etc.) together can theoretically absorb light over the entire underwater spectrum, the spatio-temporal distribution of absorbing pigments within individual cells will directly affect the absorption coefficient. The identity of particles which dominate scattering is also poorly known, largely because we understand very little concerning the composition, trophic structure, and interrelationships of plankton communities of the open ocean. One of the more significant finds of the last several years is the food web, grazers, producers, and remineralizers, composed of organisms less than 10 microns in diameter. The dynamics of this food web is largely unexplored, but it is possible that much of the flux of matter and energy in the open ocean occurs in this size range of organisms.

Related to this is the problem of detrital and organism aggregates, sometimes called "marine snow." Much of our current knowledge of the occurrence and nature of marine snow is anecdotal. Recent work suggests its importance to production and food web dynamics (e.g., Knauer et al., 1982; Caron et al., 1982; Goldman, 1984); however, quantitative data on aggregates less than 200 microns in diameter is nonexistent, photographic evidence of large (>5 mm diameter) aggregates (Fjorfi et al., 1984) suggests that they are not so abundant as to affect significantly light attenuation. Even so, aggregates, as important




Fig. 1. A conceptual model for Blowatt.

TABLE 1. Bioluminescent Taxa Responsible for Epipelagic Bioluminescence in the Sargasso Sea (0-100 m) Giving Concentrations Per Cubic Meter, C, Bioluminescence Potential, BP, and Relative Importance, I

Group	C	BP	I
Dinoflagellates			
Pyrocystis	30	$2 \times 10^{10}$	
Other	30	$10^9$	
Copepoda			
Pleuromamma	2	$2 \times 10^{12}$	1/10
Others	30?	$10^{11}$	
Ostracods	10	$1 \times 10^{11}$	6/10
Euphausiids			
Adult	0.8	$1 \times 10^{12}$	1-2/10
Larvae	8	$4 \times 10^{10}$	1/10
Decapods			
Sergestids	0.002	$10^{13}$	
Others	0.0001	$10^{15}$	
Mysids			
Amphipods	?	$10^{11}$	
Larvaceans	20	$4 \times 10^{10}$	
Coelenterates	5	$2 \times 10^{10}$	
Polychaetes	0.3	$6 \times 10^{10}$	1/10

Data courtesy of Elijah Swift, Graduate School of Oceanography, URI, Kingston, RI 02881.

absorbers and scatterers, may be missed, since current optical instruments measure a small volume and there is presently no means of extrapolating to bulk optical properties over the water column. Furthermore, it is possible that aggregates strongly influence food web dynamics. So even if their direct effect optically is small, their indirect effect could be large.

### Bioluminescence in the Ocean

A major goal of Blowatt is to predict patterns of oceanic bioluminescence and relate these to variability in optical properties over appropriate spatial and temporal scales. For purposes of modeling bioluminescence potential in the sea, an immediately obvious generalization is that the numbers and types of bioluminescent members of a pelagic community define the limits of the bioluminescent potential. That is to say, it should be possible to predict the bioluminescence potential from knowledge of the distributions of bioluminescent organisms.

Predictions of actual bioluminescence are complicated by considering that (1) light is generated in the ocean by six of the seven known bioluminescent systems; (2) in some marine environments, over 97% of the individual organisms and 90% of the species have been reported to be bioluminescent (Young, 1983); (3) spectral emission ranges in color range from the near UV to the far red; (4) kinetics range from 60 ms flashes to continuous glows lasting several minutes; and (5) intensity of the emission is controlled by a variety of biological mechanisms. Furthermore, the actual bioluminescent signal which can be elicited by strong mechanical stimulation is modulated by the physiological state of the organisms, reproductive and nutritional status, etc., as well as a variety of environmental factors.

Undoubtedly, the least understood and perhaps the most intractable aspects of bioluminescence are the biological determinants of bioluminescent response. Nevertheless, significant progress has been made in oceanographic bioluminescence research. For example, instead of dinoflagellates and fish, zooplankton have been found to be the most important bioluminescent source in the Sargasso Sea (Table 1) and perhaps in many areas in the open ocean. Recent observations point to clear relationships between temperature fronts, chlorophyll *a* distributions, and stimutable bioluminescence (Figure 2). Other observations (E. Swift et al., manuscript in preparation, 1984) show variable depth coherence between the deep chlorophyll maxima and bioluminescence potential. In the Sargasso Sea, the bioluminescence potential peak has been observed to be as much as 40 m above the chlorophyll maximum, while in some of the island passages in the Caribbean the peaks are coincident.

Spatio-temporal scales of bioluminescence are largely biologically determined. In some areas, such as the Norwegian Sea, bioluminescence is strongly seasonal because of the pop-

ulation dynamics of crustacean zooplankton. In the Sargasso Sea, on the other hand, there appears to be relatively little seasonality in numbers of relative abundance of bioluminescent organisms (E. Swift et al., manuscript in preparation). Biogeographic communities have been defined in which the relative numbers of bioluminescent and nonbioluminescent members have a stable relationship (McGowan, 1978). It is possible that population and biological studies of species responsible for much of the bioluminescence potential will be particularly useful in predicting patterns of bioluminescence.

### Field Studies

The first Blowatt experiment will be a cruise, now scheduled for April 1985. The general plan for this initial phase of Blowatt will be to sample different water masses in the northwest Atlantic for biological and optical properties and for bioluminescence potential. The questions we ask during this phase of the program are ones of composition, identification, size distributions, and pigment distributions of the biological populations. We are particularly interested in the vertical structure of physical and optical properties, biological variables, and bioluminescence potential and their interrelationships.

The overall success of the program is highly dependent on the resolution of relevant temporal and spatial scales of variability in physical, biological, and optical parameters. From a biological standpoint, the two most important time scales are the diurnal and the seasonal. The diurnal time scale is accessible from shipboard platforms; however, seasonal time series for open ocean locations are rare. We believe a surface mooring to be the best means of characterizing the temporal evolution of physical, biological, bioluminescence, and optical parameters at a given location, and this is planned for the subsequent field effort in 1987. The mooring will provide an opportunity to sample for a long enough period to evaluate seasonal cycles, with enough sampling density to assess aperiodic variability associated with synoptic weather systems. The mooring will be equipped with a wide variety of the latest physical, biological, bioluminescence and optical sensors. During cruises to service the mooring, shipboard programs can be accommodated to sample short-term variability not accessible from the in situ observations, to allow various kinds of experimental programs to help interpret mooring observations, and to characterize spatial variability and horizontal gradients in the vicinity of the mooring.

Again, the overall and long term goal of this study is to be able to make predictions of the optical properties and bioluminescence patterns of the sea from a knowledge of physical forcings, biological processes, and the interrelationships between physical, biological, and optical properties. A key link in this effort is the distribution and variance of biogenous absorbers and scatterers, their size and pigment distributions, and the dynamics of these biological populations. Clearly, this will require concerted efforts by physical, biological, and optical oceanographers.

Investigators interested in Blowatt are invited to contact the Blowatt Project Office, care of Lamont-Doherty Geological Observatory, Palisades, New York 10964.

### References

- Blizard, M. A., E. O. Hartwig, and B. J. Zahuranc, Report on the Office of Naval Research Marine Bioluminescence and Optical Variability Advisory Group Meeting, Office of Naval Res., Arlington, Va., 1982.
- Caron, D. A., R. G. Davis, L. P. Madin, and J. Sieburth, Enrichment of heterotrophic bacteria and bacteriivorous protozoa in oceanic microaggregates, *Science*, 218, 795-796, 1982.
- Dickey, T., and J. J. Simpson, The sensitivity of upper ocean structure to time varying wind direction, *Geophys. Res. Lett.*, 10, 133-136, 1983.
- Goldman, J. C., Oceanic nutrient cycles, in *Fluxes of Energy and Materials in Marine Ecosystems: Theory and Practice*, edited by M. J. R. Fasham, pp. 137-170, Plenum, New York, 1984.
- Honjo, S., K. W. Doherty, Y. C. Agrawal, and

V. L. Asper, Direct optical assessment of large amorphous aggregates (marine snow) in the deep ocean, *Deep Sea Res.*, 31, 67-76, 1984.

Klein, P., A simulation of the effects of air-sea transfer variability of the structure of marine upper layers, *J. Phys. Oceanogr.*, 10, 1824-1841, 1980.

Klein, P., and B. Coste, Effect of wind-stress variability on nutrient transport into the mixed layer, *Deep Sea Res.*, 31, 21-37, 1984.

Knauer, G. A., D. Hebel, and F. Cipriano, Marine snow: Major site of primary production in coastal waters, *Nature*, 300, 630-631, 1982.

McGowan, J. A., What regulates pelagic community structure in the Pacific, in *Oceanic Sound Scattering Prediction*, edited by N. R. Andersen and B. J. Zahuranc, pp. 423-444, Plenum, New York, 1978.


Swift, E. W., H. Biggley, P. G. Verity, and D. T. Brown, Zooplankton are major sources of epipelagic bioluminescence in the Southern Sargasso Sea, *Bull. Mar. Sci.*, 33, 855-889, 1985.

Young, R. E., Oceanic bioluminescence: An overview of general functions, *Bull. Mar. Sci.*, 33, 829-845, 1983.

This information report was contributed by John Marra of the Biological Oceanography Group at Lamont-Doherty Geological Observatory of Columbia University, Palisades, NY 10964 and Eric O. Hartwig of the Office of Naval Research, Arlington, VA 22217.

## News & Announcements

### W. Stanley Wilson: AGU Ocean Sciences Award



The Ocean Sciences Section of the AGU recognizes W. Stanley Wilson for his unique leadership contributions to the emerging role of satellite measurements in oceanography. Through his persistent efforts the ocean sciences now stand on the brink of a new era that will merge conventional research techniques with new satellite technologies.

Stan completed his B.S. and M.A. degrees at the College of William and Mary. He spent three years as a research biologist in Australia, New Zealand, and Antarctica before obtaining a Ph.D. in oceanography from The Johns Hopkins University in 1972.

He then joined the Office of Naval Research (ONR) as a program manager for physical oceanography. Promoted to program director for physical oceanography in 1976, he helped to develop and manage outstanding multi-institutional programs that included the Mid-Ocean Dynamics Experiment (MODE), the North Pacific Experiment (NOPAX), and the Mixed Layer Experiment (MILE). Stan was highly successful in integrating and translating the new research results into Navy operational use. For these efforts, he was awarded the Department of the Navy Superior Civilian Service Award in 1979.

In 1979, Stan accepted a new challenge with NASA as Chief of the Oceanic Processes Branch in the Office of Space Science and

Applications. Seasat had recently failed after only 96 days, and its successes or failures were as yet unknown. By implementing a plan for the utilization of these first-ever satellite data intended for ocean research, he and NASA have demonstrated the tremendous promise of oceanography from space.

His branch is now planning a series of new satellite missions for oceanography. These are a NASA scatterometer (for wind measurements) to fly on the Navy Remote Ocean Sensing System (NROSS) in 1989, an improved radar altimeter to measure ocean circulation for the Ocean Topography Experiment (TOPEX) in 1989, and Ocean Color Imager (OCI) for biological research. These missions have been designed through the close coordination of the NASA engineering and ocean sciences communities, as well as the Office of the Oceanographer of the Navy, the National Science Foundation, ONR, and NOAA. Stan has recognized the importance of the cooperation between different scientific disciplines. With skill, dedication, and patience he has advocated ideas through inter-agency planning. Working in concert with the scientific research, engineering, and operational oceanography communities, oceanography now has the potential for longer-term, larger-scale studies of the global oceans.

In summary, Stan's exposure to the many facets of oceanography while at ONR has served the ocean sciences well at NASA. His high energy, coupled with scientific foresight, has introduced the new satellite technologies to ocean research. He has had the tenacity and courage to break this new ground, as well as the statesmanship to convince the traditionalists in the oceanographic community of the potential value of satellite ocean remote sensing. Oceanographers are now looking to future programs that will incorporate satellite measurements as a crucial element. For his central role in the establishment of satellite remote sensing as a proven technology in ocean sciences, we recognize Stan Wilson's achievements.

This item was contributed by Christopher N. K. Moores, Past-President, Ocean Sciences Section; Joseph L. Reid, President; and Peter G. Brewer, former Secretary.

## World Ocean Circulation Experiment: Planning for a U.S. Component

A World Ocean Experiment (WOCE) to improve our description and understanding of the world ocean circulation is being planned. Technological and scientific developments of the last decades have made possible the serious consideration of such a global experiment to begin in the late 1980's. These developments include an increased understanding of the nature of ocean circulation and the sampling procedures needed to observe it, instrumentation for long time series measurements, numerical ocean models and high-capacity computers to use them, improved methods for measurement of chemical tracers, the means for obtaining improved ocean measurements from vessels of opportunity, satellite technology that can observe both the primary atmospheric forcing and the oceanic response, and the realization that addressing global societal problems related to the ocean will require a coordinated, global-scale program of ocean observation and modeling.

The WOCE is considered to be a contribution to the study of long-term climate trends and sensitivity (Stream 3) of the World Climate Research Program (WCRP). At the international level the formulation is being guided by a Scientific Steering Group under the auspices of the Committee on Climatic Changes and the Ocean and the Joint Scientific Committee of the WCRP.

The provisional goals for an international WOCE are (1) to collect the data necessary to develop and test ocean models for predicting climate change and (2) to determine the representativeness of the specific WOCE data sets for the long-term behavior of the ocean and find methods for determining long-term changes in the ocean circulation. Since the focus is on the construction of ocean models and the collection of data sets necessary for demonstrating that these are useful models of the ocean circulation, the determination of what these data sets should be is the crux of the WOCE design. The International Scientific Steering Group has established a Numerical Experimentation Group and six working groups in different areas to consider what major data sets are required and to determine strategies for obtaining the required data.

On the basis of present concepts it seems likely that the major elements of WOCE will include models, satellite altimeters, a satellite gravity mission, satellite scatterometers for surface winds, other fields of surface forcing, hydrography, tracer measurements, direct velocity measurements, acoustic tomographic arrays, ship-of-opportunity programs, and a data management system. Many of these elements must be coordinated with the proposed



## Oceanography (cont. from p. 733)

satellite missions scheduled for the 3- to 5-year lifetimes to begin in 1989. Others must begin earlier (now) and continue throughout the most intensive operational phase and beyond.

Each participating country is expected to develop a national plan which might contribute in the areas of its capability and interest to the overall experiment. Within the United States this planning process began with a workshop on "Global observation and understanding of the general circulation of the oceans" for some 60 members of the U.S. oceanographic community. The workshop was organized under the auspices of the National Research Council (NRC) Board on Ocean Science and Policy and held during August 1983 in Woods Hole, Mass. The workshop participants agreed that the WOCE concept is worthwhile and timely, identified a tentative U.S. goal and objectives, and recommended that a U.S. planning committee and a number of working groups be established to address critical issues. The provisional objectives of the U.S. component of WOCE are directed toward describing and understanding global ocean circulation. It was felt that this understanding is important for itself and must precede an understanding of the role of the ocean in climate.

The workshop report received wide distribution and review, and a panel for the U.S. WOCE was constituted within the NRC, sponsored jointly by the panels of the Board on Ocean Science and Policy and the Board on Atmospheric Sciences and Climate. This panel has been moving toward the design of a U.S. WOCE component. Initial support for the panel activities has been provided via the NRC and by a grant from the National Science Foundation to the Joint Oceanographic Institutions Inc. Several U.S. working groups (closely coordinated with the international groups) have been established: communication links between the U.S. scientific community, the sponsoring agencies, and the international components are being implemented; and a proposal for an intensive planning activity for the fiscal years 1985-1987 has been prepared.

To date, U.S. working groups have been formed to plan WOCE activities in the following areas: measurement and interpretation of tracers, experimental design for measuring geostrophic circulation, surface forcing (wind stress and heat and moisture fluxes), numerical modeling, data management, and technology development. In addition, other ad hoc group meetings and activities by interested groups of scientists are being supported.

To inform the community of the activities of the panel, working groups, and ad hoc groups, we plan to publish, in TOR, brief reports of meetings and other items of interest. In this issue we include the report of a Deep Drifters Meeting held in Denver, Colo., on May 18-19, 1984. Interested scientists should contact panel members, meeting chairpersons, working group members, or the U.S. WOCE Planning Office: W. Nowlin, Department of Oceanography, Texas A & M University, College Station, TX 77843 (telephone: 409-845-2947; telemail: Sciencenet, W. Nowlin).

## New Research Vessels

Two "new" ocean-going research vessels operated by the Scripps Institution of Oceanography and the National Science Founda-

tion (NSF) will soon begin full-time scientific duties off the coast of California and in the Antarctic, respectively. The 37.5-m Scripps vessel, named *Robert Gordon Sprout* in honor of the ex-president of the University of California, replaces the smaller ship *Ellen B. Scripps*, which had served the institution since 1965. The new ship is a slightly modified Gulf Coast workboat. Under the name of *Midnight Alaskan*, it had been used for high-resolution geophysical surveys in American and Latin American waters by such firms as Arco Oil & Gas, Exxon, Pennzoil, and Rascal-Decca before its purchase by Scripps from a Louisiana chartering firm last summer.

The *Robert Sprout* is undergoing a number of modifications for scientific outfitting, including the addition of laboratories, winches, booms, and electronic and research instruments. The ship will be used mainly along the California coast and in the Gulf of California for biological investigations, physical oceanography, and scientific equipment testing. Its maiden voyage, however, scheduled for early September, will take it to Yucatan, Mexico, for a study of seals, then on to the west coast of Mexico for in-transit studies of marine mammals and birds. The *Robert Sprout* is expected to return to its home port of San Diego by late October, and to make its first voyage along the California coast in support of physical oceanography experiments in late autumn. Meanwhile, the *Ellen B. Scripps* is being offered for sale, with the proceeds to be used to pay for the new ship.

The new NSF Antarctic research ship, *Polar Duke*, will replace the wooden-hulled *Hero*, which has served in the Antarctic since 1968. The new ship is larger (85.7 m versus 37.5 m), and its hull is classified as strong as that of an ice-breaker. It will have room for approximately 40 people (some 2-3 times *Hero's* capacity), will carry more laboratory equipment, and will have a helicopter deck. As a result, it will be able to conduct research in the rough waters between South America and Antarctica that was impossible with the aging *Hero*. The new ship's cargo space is being converted to laboratories in preparation for its first scientific voyage on or about January 1 of next year. The *Polar Duke* will support antarctic research in biology, oceanography, and geology. Originally designed for scientific and transport expeditions, the vessel is being leased from Garmco Shipping Limited of Newfoundland by ITT/Antarctic Services, who will operate the ship for the NSF for a period of 3 years. There is also a provision for two 1-year extensions. *Hero*, meanwhile, will be retired after 16 years of operations in Antarctic waters, and current plans are to dispose of it as government surplus.

## Meetings

## Deep Drifters Meeting

Technical development of a new generation of deep drifting systems is proceeding at several locations. A meeting was held in Denver, Colo., on May 18-19, 1984, under the joint sponsorship of the National Research Council (NRC) Panel for a U.S. World Ocean Circulation Experiment (WOCE) and the Drifters development program. The purpose was to discuss the state of development of these systems, their prospects for continued development, and their possible use in programs such as WOCE, where the intended spatial and temporal sampling scales are quite

large. The three new systems discussed were RAFOS (SOFAR spelled backwards), the General Circulation Drifter, and Dyogene. The capabilities of existing Sound Fixing and Ranging (SOFAR) floats were also discussed.

The expectation of remotely sensed data from satellites as part of the WOCE has prompted consideration of what types of in situ instruments are most suitable for use with such data. Satellite data approaches global spatial coverage and continues over years, with a temporal resolution between a few days and a month. Very few in situ measurement systems can approach these sampling characteristics without heroic effort and extraordinary expense. Drifting buoy systems, which report sensor data and/or position through satellite systems, such as the present System ARGOS, hold great promise for achieving sampling characteristics which are compatible with satellite data if they can be made sufficiently inexpensive to be deployable in large numbers. The focus of this meeting was on drifting buoys capable of measuring interior ocean currents by passive drift.

A major objective of WOCE is to determine the transports of heat, salt, and other chemical properties over large distances due to both systematic (mean) and irregular (diffusive) advection. Traditionally, these transports have been inferred largely from the spatial distributions and knowledge of sources and sinks. These inferences can be made in principle with more confidence for transient property fields (e.g., bomb radiocarbon and tritium). However, as a complement to such indirect methods, the long-time trajectories of deep drifting buoys may provide important direct information about the capacity of the general circulation for transport of passive water properties.

The Global Circulation Drifter (GCD) is under development by Doug Webb at Webb Research Corporation in collaboration with Russ Davis at Scripps Institution of Oceanography. It is designed as a free-drifting vehicle which will operate down to 2000 m and be capable of several round trips to the surface to report its position by ARGOS. The design is for a lifetime of 10 years and of approximately 40 round trips to the surface, which might be extended to 200 with further development. The GCD is undergoing testing at sea in 1984.

RAFOs is a buoy which drifts at the depths of the sound channel. It is under development by Tom Rossby at the University of Rhode Island. In addition to temperature and pressure, RAFOs stores the time of arrival of signals from fixed SOFAR sound sources three times a day. At the end of its design lifetime (presently 6 months) the float surfaces and telemeters its stored data through ARGOS. The sound sources are moored in the area of interest on simple moorings. The usable range from source to the RAFOs float is about 1500 km, though this will vary geographically depending on the water characteristics. Acoustic sources have an expected lifetime of perhaps 2.5 years, which might be extended to 5 years with additional development. Similarly, the lifetime of the RAFOs floats themselves probably can be extended to beyond a year. This system will be used in the Gulf Stream and adjacent regions during the period 1984-1986.

Dyogene is under development by J. C. Gascard in France. It is a deep drifter designed to sample for periods of about 1000 days, with data subsets of about 100 days being reported via ARGOS during a round trip to the surface. Dyogene rises by releasing ballast and descends by releasing buoyancy in the form of glass balls. A 1-month test on the surface confirmed the performance of the data-reporting system, including a low-profile antenna. Depth cycling tests are planned next. A pilot experiment with a limited number of buoys will be carried out in late 1988 or 1987. The lifetime of Dyogene is presently limited by the life of the lithium batteries. It could be extended by using solar rechargeable cells.

SOFAR floats have been used in a variety of locations and experiments since the late 1960s. In its original form the system consisted of drifting sources and shore-based receivers. Later innovations include the use of a moored recoverable receiver, the Autonomous Listening Station (ALS), which frees the experimenter from the geographical constraints of working near existing shore stations. An even more recent development is the RELAYS (Real-Time Link and Acquisition Yare) system, which may be regarded as a drifting ALS. This system is being tested during 1984. The present expected lifetime of SOFAR floats is about 2.5 years, which can be expected to be increased with better quality control but minimal redesign. About 50% of the floats deployed have lasted to their energy limit (battery life). SOFAR floats may drift at various depths in the range 500-2000 m, with the best performance (greatest range) obtained in the sound channel.

Contemplating the use of drifters in large-scale experiments as part of the WOCE leads to the question of operational feasibility. Experimental design may suggest the use of 1000 floats in an ocean basin. How are such large numbers of floats to be prepared and deployed? Will it be necessary to establish new facilities to carry out this operational ac-

tivity and relieve the burden on the more experiment-oriented academic institutions? Such facilities might handle the drifting floats, moored sources, and moored Autonomous Listening Stations.

The ARGOS transmitter is a substantial part of the cost of construction of a drifting buoy. Less expensive ARGOS-approved Position Transmittal Terminals (PTT) with position accuracy of about 5 km (lower than present) may be commercially obtainable. An accuracy of 5-10 km would be adequate for the buoys under discussion, and a lower-cost PTT should be sought and approved for ARGOS use.

The discussion of a plan for deep drifters in WOCE addressed two general questions: For what purposes would one want the deep drifter data? How many drifters would be needed? The proposed purposes were analyses of mean horizontal velocity and velocity variance (eddy kinetic energy), dispersion of drifters (hence horizontal diffusivity), mapping of the very large scale and low frequency velocity field, and pathways of the general circulation, i.e., displacements over distances larger than individual eddy scales. For each purpose one can determine how many drifters are required to achieve a given degree of success.

These determinations have, as yet, been made only very crudely. However, there was general agreement that something near 1000 drifters years of data in an ocean basin the size of the North Atlantic would make a significant contribution to each of these purposes. (This assumes moderately frequent location reporting, which is more easily accomplished for acoustically tracked drifters, and moderately uniform horizontal distribution.) A crude illustration of the basis for this number, most relevant to the first of the purposes above, follows. One thousand drifters, reporting weekly for a year and uniformly distributed in an area equal to a 4000-km square, would give 50 nearly independent samples of low-frequency velocity on a grid spacing of about 125 km. This would determine the mean with an accuracy better than 1 cm/s for a standard deviation of 5 cm/s. If the analysis grid were coarse, the accuracy would be better for the same number of drifters, or the same accuracy would be achieved with fewer drifters.

The following points relevant to future development of buoy systems were made:

- The RAFOs receiver should be put in the GCD (or Dyogene) vehicle, providing a multiply recycling, acoustically traced drifter.
- These floats should be aimed at intermediate depths. The flow at extreme depths is more topographically controlled and therefore harder to interpret in a general circulation context, while the shallower depths (above 500 m) can probably be measured with cheaper drogued surface buoys. (SOFAR and RAFOs are inherently restricted to intermediate depths in any case, because of the need for reasonably long acoustic ranges.)
- Effort should be put into the technical aspects of combining RAFOs, SOFAR, and ocean acoustic tomography in future experiments, using a mutually compatible system of acoustic sources.

• Consideration should be given to adding data telemetry to the SOFAR Autonomous Listening Stations to allow "real-time" processing of data. The Moored Oceanographic Instrumentation System (MOIST) program at Woods Hole Oceanographic Institution might provide the technology base to make this possible.

• Both slips-of-opportunity and aircraft should be considered for buoy deployment, and each buoy system should be so planned. This new generation of intermediate depth drifting floats will provide new tools for the general circulation investigator. Their capabilities are such that a selection or mix of float types may be called for in a given experiment, depending on experimental design and location. There are intriguing possibilities for synergistic combinations between varieties of floats and between the floats and other new technology, such as tomography.

## Acknowledgments

Funding for the WOCE planning activity was provided by the National Science Foundation; support for the Drifters program has been provided by the National Oceanic and Atmospheric Administration, the National Aeronautics and Space Administration and the Office of Naval Research.

This report was submitted by Robert Heinmiller of Onnet, Inc. and James McWilliams of the National Center for Atmospheric Research (NCAR).

## ATTENTION SUBSCRIBERS!

Beginning in 1985:  
Reviews of Geophysics  
and Space Physics  
will be titled  
Reviews of Geophysics.  
Approximately 600 pages  
to be published in  
Volume 23, 1985.

## News (cont. from p. 729)

eruption on the southern part of the fissure had ceased except on one crater, which then changed into phreatic activity.

"On the northernmost part of the fissure, activity continued as of the morning of 12 September with significant lava production. By 8 September, inflation rates over the magma chambers diminished and slow deflation started."

"As in previous eruptions, lava production was highest on the northern part of the fissure and has so far not constituted any threat to inhabited areas."

Information Contacts: Karl Grönqvist, Nordic Volcanological Institute, University of Iceland, Reykjavik, Iceland; Páll Einarsson, Science Institute, University of Iceland, Reykjavik, Iceland.

## Meteoritic Events

Fireballs: Canada; Czechoslovakia; Netherlands; Tasman Sea; California, south-central states, Massachusetts, Oregon, Wisconsin, USA; Zimbabwe.

## Books

## Groundwater Contamination

Panel on Groundwater Contamination of the Geophysics Study Committee of the National Research Council, National Academy Press, Washington, D. C., xii + 179 pp., 1984.

Reviewed by R. Allan Freeze

Since 1977, the National Research Council (NRC) has published 18 monographs in its *Studies in Geophysics* series. Earlier volumes have dealt with such diverse topics as energy and climate, geophysical prediction, continental tectonics, and explosive volcanism. *Groundwater Contamination* is the 14th volume in the series. It was produced by a Panel on Groundwater Contamination initiated by the Geophysics Study Committee of NRC in consultation with representatives of the supporting agencies and members of the scientific community. The preliminary scientific findings formed the basis of an American Geophysical Union symposium in San Francisco in December 1981. This monograph is the most concise and most accessible reference available on this important environmental problem.

The report consists of 14 chapters authored by well-known researchers in the field and an overview of the study that summarizes the highlights of the chapters and formulates conclusions and recommendations. The 14 chapters are arranged in five groups: 1, Background (one chapter); 2, Processes (two chapters); 3, Methods of Waste Disposal (two chapters); 4, Examples (seven chapters); and 5, Institutional Aspects (two chapters).

The Background chapter by Veronica Pye and Jocelyn Kelley of the Academy of Natural Sciences of Philadelphia assesses the extent of groundwater contamination in the United States on the basis of two independent regional surveys that have been carried out, one by the Environmental Protection Agency and one by the Environmental Assessment Council. They conclude that cases of groundwater contamination have been documented in all parts of the country but that "we are still in a position to make choices on how best to use, manage, and protect this valuable resource."

The chapters on Processes by Mary Anderson of the University of Wisconsin and John Cherry and his coworkers at the University of Waterloo provide an up-to-date assessment of our physical and chemical understanding of the movement of contaminants in groundwater. While the current advection/dispersion/retardation approach is capable of mimicking observed behavior, it is now clear that it does not represent an understanding of the transport processes at the fundamental level required. There is still much to be learned with respect to organic contaminants, transport in fractured rock, and the questions of scale that surround the slippery concept of dispersion.

To my mind, the heart of this book lies in the seven case histories presented in part 4. It is in the individual tales of mismanagement, realization, and reclamation that the difficulties in measurement, assessment, and prediction become clearest. Many of the most visible of the well-known recent contamination incidents are here: The Love Canal in New York, the Rocky Mountain Arsenal in Colorado, the Llangollen landfill in Delaware, and the low-level nuclear waste dumps at Hanford, Washington, and Maxey Flats, Kentucky. The description of the aquifer reclamation project at the Rocky Mountain Arsenal by Leonard Konikow of the U.S. Geological Survey and Douglas Thompson of the U.S. Army Corps of Engineers is a particularly valuable case history.

In the Overview prepared by the panel, there are four recommendations that appear in italics:

1. Research is needed on the effects of chemical reactions on transport and dispersion of contaminants by groundwater and the quantification of flow in fractured media.
2. There should be a more thorough search for disposal sites that can be used safely to isolate toxic wastes from the biosphere for long periods.
3. A strategy [should] be developed that provides for the segregation, treatment, and disposal of wastes according to their hazards and their chemical affinities.
4. Governmental and industrial organiza-

tions need to agree whether various classes of wastes should be disposed of locally or in regionally designated repositories.

This is a valuable book. It will undoubtedly be used for readings in many hydrogeology seminars, and it is well suited to this purpose. But it is even more important that its message reach environmental engineers, designers of waste disposal systems, administrators of local and state agencies, and lawyers and legislators.

R. Allan Freeze is a professor in the Department of Geological Sciences at the University of British Columbia, Vancouver, Canada.

## Physical Geology: Principles and Perspectives

E. A. Hay and A. L. McAlester, Prentice-Hall, Englewood Cliffs, N.J., 1984.

Reviewed by James W. Shehan, S.J.

Textbooks on physical geology have proliferated over the past 20 years or more, during which time most fields have undergone a subject matter explosion. The challenge of authoring such a textbook is to accurately summarize the most important factual and theoretical results and to present the material in a pedagogically attractive and meaningful manner. The authors mainly have met that challenge.

Basically, I could teach the course quite happily using this book as the text. An attractive feature is that the authors have summarized those aspects of the science with which I am most familiar in a generally acceptable and interesting manner. This includes excellent line drawings and block diagrams calculated to be very helpful to the student. Those chapters dealing with areas of the authors' scientific expertise are understandably stronger and smoother than some others. A useful perspective for the beginning student and teacher is an explicit discussion of geology as a science with a comparison and contrast of methods and results in relation to other fields of science.

The chapters are brief enough to provide a readable overview. Following the treatment of classification and principles, a number of chapters have several pages illustrating the foregoing by an analysis of regional geological features of considerable general interest, such as Yosemite, Kings Canyon, and Sequoia National Parks noted for igneous activity, for example. Each chapter has an abbreviated, useful outline of the main points covered and several appropriate references for additional reading. Many chapters have interesting, short single topic treatments ("boxes") (e.g., "The Driving Mechanism Behind Plate Tectonics"). The book is a good teaching vehicle because, with notable exceptions, it is easy to supplement the text in class with one's own favorite illustrations. Such an exception is chapter 2, "Basic Building Blocks—Matter and Minerals," which would be more effective if abbreviated, and some of the exquisite detail of crystal chemistry stored in an appendix for one who needs a fuller explanation.

There are a few incomplete sentences, inexact or awkward usage of terms such as silica for siliceous, or "ultramafic igneous rocks and serpentine" (p. 285), omission of sheeted dikes in description of ophiolites, or neglect of the solar wind in the consideration of the shape of the lines of force of the earth's magnetic field (p. 259, Figure 7.10), for example. These are all mainly items that proofreaders and paid consultants will remedy for the second printing of this revised edition, no doubt. A number of the photographs (black and white) tend to be too dark or are otherwise of inferior quality (e.g., Figure 4.25b) and detract from the generally outstanding quality of the line drawings.

This is an attractive book mainly because it is a carefully abbreviated summary of most of the topics and is very well illustrated, notably with line drawings. The pedagogically useful references to the relationships between geology and the needs of mankind, chapter summaries, vignettes of regional geology classics, "boxes," and additional readings, and a good quality glossary and index outweigh the negative points noted above. In general, such features of this revised edition should render it

## Earthquakes

Date	Time, UT	Magnitude	Latitude	Longitude	Depth of Focus, km	Region
Aug. 6	1907	6.8 M <sub>s</sub>	32.40°N	131.81°E	50	Off Kyushu, Japan
Aug. 17	1806	3.9 m <sub>LG</sub>	37.82°N	78.39°W	6	SW Virginia, USA

Information Contacts: National Earthquake Information Service, U.S. Geological Survey, Stop 967, Denver Federal Center, Box 25040, Denver, CO 80225.

quite competitive with some of the other books of quality now available.

James W. Shehan, S.J., is director of Weston Observatory and professor of geology, Department of Geology and Geophysics, Boston College, Weston, MA 02193.

## Nonlinear Waves

Lokenath Debnath (Ed.), Press Syndicate of the University of Cambridge, New York, 360 pp., 1983.

Reviewed by Dennis Papadopoulos


The book *Nonlinear Waves* edited by L. Debnath is a collection of papers on the theory and applications of nonlinear wave phenomena. The book is an outgrowth of an NSF-CBMS regional research conference on nonlinear waves and integrable systems, convened in June, 1982 at East Carolina University. The book aims at bringing together recent developments on the theory of nonlinear waves and solitons, with selective applications in the areas of fluid dynamics and plasma physics. The field had an explosive growth during the last decade, and it is, correctly, expected that it will exert a major influence on future research directions of many fields of physics and engineering.

The book is divided into three parts. The first two deal primarily with applications of the mathematical techniques for solving nonlinear equations to special problems in the fields of fluid dynamics and plasma physics. The third part deals mainly with applied

mathematics and focuses on recent results and extensions of the inverse scattering transformation (IST), of the evolution equations, and of the statistical mechanics of the Sine-Gordon Hamiltonian. A total of 18 papers are included in the volume, seven on fluid dynamics, five on waves and solitons, and six on applied mathematics. Taken individually, almost all of the papers are well written reviews of important subjects in their specific fields, including a variety of recent and exciting results and techniques. Of particular interest are the fluid mechanics chapters 2-5, along with chapters 16 and 17 of the applied mathematics part, which discuss a variety of solutions of the Korteweg-deVries (KdV) and nonlinear Schrödinger (NLS) equations for systems with and without dissipation, including the role of variable coefficients. A general nth order spectral transform and its inversion and the role of recurrence are also presented in an interesting fashion.

Unfortunately, that is as far as I can point to any kind of unifying thread among the various chapters of the book, and even in this case I am stressing it. The level of mathematical sophistication is so uneven as to make most of the book rather unreadable on any level of general physics, applied mathematics, or engineering audience. I do not believe that the desired cross fertilization among the nonlinear hydrodynamicists, plasma physicists, and applied mathematicians can be accomplished with this book. It is possible that this could have been achieved through the discussions at the conference; nevertheless, as often happens in similar cases, the follow-up manuscripts tend to be more sophisticated and spe-

Books (cont. on p. 731)



This outstanding series assesses geophysical knowledge in specific fields, particularly as it relates to societal problems. The series is a cooperative undertaking of the National Research Council's Geophysics Research Forum and the American Geophysical Union.

**Groundwater Contamination**  
John D. Bredehoeft, Panel Chairman  
Using both hydrologic theory and data gained from the study of contaminated aquifers, this new volume explores current knowledge of how contaminants move in groundwater. The report stresses the need for a waste-disposal strategy for the prevention of future groundwater contamination. 1984, 179 pages, paperback, \$17.95

**Explosive Volcanism: Inception, Evolution, and Hazards**  
Francis R. Boyd, Jr., Chairman  
Papers by well-known scientists present sound technical information on various aspects of explosive volcanism: mantle chemistry and processes, tectonic influences, subduction geometry, volcanic periodicity, mechanisms of magma formation and segregation, and eruption mechanics. The authors underscore the interrelationships among geophysical events and recommend directions for future research and monitoring efforts. 1983, 192 pages, 3 maps, clothbound, \$24.50

**Other volumes in the series...**  
*Fundamental Research on Estuaries: The Importance of an Interdisciplinary Approach*; C. B. Officer and L. E. Cronin, Cochairmen, 1983, 79 pp., \$6.95 • *Climate in Earth History*; W. H. Berger and J. C. Crowell, Cochairmen, 1982, 198 pp., \$16.25 • *Solar Variability, Weather, and Climate*; J. A. Eddy, Chairman, 1982, 106 pp., \$11.95 • *Scientific Basis of Water-Resource Management*; M. B. Fiering, Chairman, 1982, 127 pp., \$12.25 • *Mineral Resources: Genetic Understanding for Practical Applications*; P. B. Barton, Jr., Chairman, 1981, 119 pp., \$12.95 • *Continental Tectonics*; B. C. Burchfiel, J. R. Oliver, and L. T. Silver, Cochairmen, 1980, 397 pp., \$14.95 • *Impact of Technology on Geophysics*; H. E. Newell, Chairman, 1979, 121 pp., \$11.75 • *Geophysical Prediction*; H. R. Landsberg, Chairman, 1978, 215 pp., \$14.25 • *Estuaries, Geophysics, and the Environment*; C. B. Officer, Chairman, 1977, 127 pp., \$9.95 • *Climate, Climatic Change, and Water Supply*; J. Wallis, Chairman, 1977, 132 pp., \$9.25 • *The Upper Atmosphere and Magnetosphere*; E. S. Johnson, Chairman, 1977, 168 pp., \$11.50 • *Energy and Climate*; R. R. Revelle, Chairman, 1977, 158 pp. (Inquire for price)  
To order these reports, please send payment to National Academy Press, Dept. JH 726, 2101 Constitution Avenue, NW, Washington, DC 20418.

# Groundwater Transport:

## Handbook of Mathematical Models (1984)

Water Resources Monograph Series 10

I. Javandel, C. Doughty, and C. F. Tsang,  
240 pages • softbound • illustrations

\$16

Written in a clear style, this work reviews, selects, and demonstrates the best and most practical mathematical models to predict the extent of groundwater subsurface contamination. Primary emphasis is given to the use of simple formulas and comprehensive tables, resulting in a readily usable guide in the field.

30% Discount to AGU Members

Call: 800-424-2488  
(202) 462-6003 (toll-free DC area or outside contiguous USA)

Write: American Geophysical Union  
2000 Morkin Ave., N.W.  
Washington, D.C. 20000

Orders under \$50 must be prepaid

accept

501984



## Books (cont. from p. 735)

cialized, breaking the unifying tie achieved through the usually simple physical presentations in the research conference.

The subject of nonlinear dynamics, including applications to physics and engineering, has made tremendous strides during the last decade. The sooner the general scientific public becomes familiar with it, the sooner we will have major breakthroughs in the understanding of many subjects in physics and engineering. Several excellent books with good descriptions of both the fundamental mathematical theory of IST and spectral methods as well as applications to fluid dynamics, plasma physics, and solid state have appeared recently. Ablowitz and Segur, "Solitons and the Inverse Scattering Transform," and Lamb, "Elements of Soliton Theory," are examples of such books, although they stress the subject of conservative systems. The role of dissipation and of strange attractors is adequately treated in *Topics in Nonlinear Dynamics*, edited by S. Jorna, as well as the proceedings of the various "Twente" Conferences. I believe that referring to the above references, especially the Ablowitz and Segur volume, can much better accomplish the goals set by the editor of the book under review here. Meanwhile, *Nonlinear Waves* can remain a collection of good review articles on the subject of nonlinear waves, but with only the title as the unifying entry.

Dennis Papadopoulos is with the Department of Physics, University of Maryland, College Park, MD 20742.

## New Publications

Items listed in New Publications can be ordered directly from the publisher; they are not available through AGU.

**Acid Rain: A Water Resources Issue for the 80's**, R. Herrmann and A.J. Johnson (Eds.), American Symposium on Hydrometeorology, iii + 83 pp., 1983.

**Aoustic Vanequid: Applications to Oceanic Science**, C.A. Boyles, John Wiley, New York, xi + 321 pp., 1984, \$46.95.

**The Correlation of Climate and Life**, S.H. Schneider and R. Londer, Sierra Club Books, \$25.

**Deconvolution**, A. Zolzkowski, International Human Resources Development, Boston, Mass., xi + 175 pp., 1984, \$36.

**Gas Transfer at Water Surfaces**, W. Brutsaert and G.H. Jirka (Eds.), D. Reidel, Hingham, Mass., x + 639 pp., 1984, \$78.

**Heat Conduction**, U. Grullig and H. Sander, (Transl. by J. Kestin), Springer-Verlag, xx + 187 pp., 1984.

**Helium Isotopes in Nature**, B.A. Mamyrin and L.N. Tolstikhin, *Dev. in Geochem.*, vol. 3, Elsevier, New York, xiv + 273 pp., 1984, \$53.75.

**International Symposium on Hydrometeorology**, A.I. Johnson and R.A. Clark (Eds.), American Water Resources Association, Bethesda, Md., ix + 598 pp., 1983.

**Kimberlites, I. Kimberlites and Related Rocks**, J. Kornprobst (Ed.), *Dev. in Petrol.*, 11A, Elsevier, New York, xiv + 408 pp., 1984, \$53.75.

**Kimberlites, II. The Mantle and Crust-Mantle Relationships**, J. Kornprobst (Ed.), *Dev. in Petrol.*, 11A, Elsevier, New York, xiv + 393 pp., 1984, \$50.

**The Major Biogeochemical Cycles and Their Interactions**, B. Bolin and R.B. Cook (Eds.), John Wiley, New York, xxi + 532 pp., 1983.

**The Morphostructure of the Atlantic Ocean Floor: Its Development in the Mesozoic**, V.M. Litvin, (Transl. by V.M. David, N.N. Protchenko and V.U. Rodzhabov), D. Reidel, Hingham, Mass., ix + 172 pp., 1984, DFL 05.

**The Near-Earth and Interplanetary Plasma**, vol. 1: *General Properties and Fundamental Theory*, V. L. Alpert, Cambridge University, New York, Melbourne, viii + 209 pp., 1983, \$59.

**The New Science of Strong Materials**, J.E. Gordon, 2nd ed., Princeton University, Princeton, N.J., 287 pp., 1984, \$9.95.

**Nitrogen in the Marine Environment**, E. J. Carpenter and D. G. Capone (Eds.), Academic, New York, xvii + 900 pp., 1983, \$59.

**Noble Gas Geochemistry**, M. Ozima and F. A. Podosek, Cambridge Univ., New York, vii + 365 pp., 1983, \$79.50.

**Nonlinear Waves**, L. Debnath (ed.), Cambridge University, New York, 390 pp., 1983, \$39.50.

**Ophiolites: Oceanic Tectonics and Metamorphism**, Suppl. 8, Ofiolita, Pitagora, Bologna, 73 pp., 1983.

**Our Threatened Climate: Ways of Averting the Problem Through Rational Energy Use**, W. Bach (Transl. by J. Jäger, D. Reidel, Hingham, Mass., xxiv + 368 pp., 1984, \$22).

**Palaeoclimatic Research and Models**, A. Ghazi (Ed.), D. Reidel, Hingham, Mass., vii + 205 pp., 1983, \$34.50.

**Permafrost: Fourth International Conference Proceedings**, July 17-22, 1983, Organized by University of Alaska and National Academy of Sciences, National Academy Press, Washington, D.C., xxv + 1524 pp., 1983.

**Phosphate Minerals**, J.O. Nriagu and P.B. Moore (Eds.), Springer-Verlag, New York, vi + 442 pp., 1984, \$57.50.

**Physical Geography: A Landscape Appreciation**, T. L. McKnight, Prentice-Hall, Englewood, NJ, x + 488 pp., 1984.

**Physical Geography: Principles and Perspectives**, 2nd ed., E. A. Hay and A. L. McLeister, Prentice-Hall, Englewood Cliffs, NJ, xii + 463 pp., 1984, \$25.95.

**Physical Oceanography of Coastal Waters**, K. F. Bowden, Ellis Horwood Ser. in Mar. Sci., New York, 302 pp., 1983, \$68.95.

**The Polarization Method of Seismic Exploration**, E.I. Galperin, D. Reidel, Hingham, Mass., xiv + 268 pp., 1984, \$58.

**Precambrian of South India**, S.M. Nayvi and J.W. Rogers (Eds.), Mem. J. Geological Society of India, Bangalore, xx + 575 pp., 1983.

**Principles of Lake Sedimentology**, L. Håkanson and M. Jansson, Springer-Verlag, New York, x + 316 pp., 1983, \$39.

**Problems and Prospects in Logging and Medium Range Water Forecasting**, D.M. Burridge and E. Källén (Eds.), Topics in Atmos. and Oceanogr. Sci., Springer-Verlag, New York, vi + 274 pp., 1984, \$17.50.

**Proceedings of the Biomas Colloquium in 1982: Special Issue No. 27**, T. Nemoto and T. Matsuda (Eds.), Mem. of Inst. of Polar Res., Spec. Issue 27, National Institute of Polar Research, Tokyo, iv + 247 pp., 1983.

**Proceedings of the Third Symposium on Antarctic Geosciences**, 1982: *Special Issue No. 28*, T. Nagata (Ed.), National Institute of Polar Research, Tokyo, iv + 289 pp., 1983.

**Proterozoic Geology**, L. G. Medaris, Jr., C. W. Byers, D. M. Mickelson, and W. C. Shanks (Eds.), Mem. 161, Boulder, Colo., viii + 315 pp., 1983.

**Regional Trends in the Geology of the Appalachians-Caledonides-Hercynides-Mauritania Orogen**, Paul E. Schenk (Ed.), NATO ASI Ser., Ser. C: Math and Phys. Sci., vol. 116, D. Reidel, Hingham, Mass., xiii + 398 pp., 1983, \$53.50.

**Remote Assessment of Ocean Color for Interpretation of Satellite Visible Imagery**, H. R. Gordon and A. Y. Morel, Springer-Verlag, New York, 114 pp., 1983, \$20.

**Revolution in the Earth Sciences: Advances in the Past Half-Century**, S. J. Boardman (Ed.), Kendall/Hunt, Dubuque, Iowa, viii + 385 pp., 1983.

**Rock and Mineral Magnetism**, W. O. Reilly, Blackie, Chapman and Hall, New York, xi + 220 pp., 1984, \$39.95.

**Rock Fracture Mechanics**, No. 275, H. P. Rossmanith (Ed.), Springer-Verlag, New York, xii + 484 pp., 1983, \$34.

**The Role of Heat in the Development of Energy and Mineral Resources in the Northern Basin and Range Province**, Special Report No. 13, Geothermal Resources Council, v + 384 pp., 1983, \$30.

**Scale-Up of Water and Wastewater Treatment Processes**, N.W. Schmidke and D. W. Smith (Eds.), Butterworth, Boston, pp., 1983, \$39.95.

**Science and Creationism: A View from the National Academy of Sciences**, Committee on Science and Creationism, National Academy of Sciences, National Academy Press, Washington, D.C., 28 pp., 1984, Quantity: 1; \$4; 2-9; \$3; 10 or more, \$1.75.

**Sediment Diagenesis**, A. Parker and B. W. Sellwood, (eds.), D. Reidel, Boston, vii + 427 pp., 1983, \$53.50.

**Seismic Design Technology for Breeder Reactor Structures**, vol. 1, *Special Topics in Earthquake Ground Motion*, D. P. Reddy (Ed.), Agabian Associates, El Segundo, Calif., ix + 267 pp., 1983, \$23.50.

**Seismic Design Technology for Breeder Reactor Structures**, vol. 2, *Special Topics in Soil Structure Interaction Analysis*, D. P. Reddy (Ed.), Agabian Associates, El Segundo, Calif., vii + 159 pp., 1983, \$16.

**Seismic Design Technology for Breeder Reactor Structures**, vol. 3, *Special Topics in Piping and Equipment*, D. P. Reddy (Ed.), Agabian Associates, El Segundo, Calif., vii + 307 pp., 1983, \$19.

**The Technology of High-Level Nuclear Waste Disposal**, vol. 1, P. L. Hoffman and J. J. Breslin, (eds.), Technical Information Center, U.S. Department of Energy, Oak Ridge, Tenn., xii + 395 pp., 1981, \$18.

**Theoretical Glaciology: Material Science of Ice and the Mechanics of Glaciers and Ice Sheets**, K. Hutter, Math. Approaches to Geophys., D. Reidel, Hingham, Mass., xxxii + 510 pp., 1983, \$104.

**Treatment Plant Hydraulics for Environmental Engineers**, L. D. Benfield, J. F. Jenkins, Jr. and A. D. Parr, Prentice-Hall, Englewood Cliffs, NJ, 231 pp., 1984, \$39.95.

**Tsunami: Their Science and Engineering**, K. Lida and T. Iwasaki (Eds.), *Ad. in Earth and Planet. Sci.*, D. Reidel, Boston, xiv + 563 pp., 1983, \$113.

**United States Geological Survey, 1981**, C.W. Stover (Ed.), U.S. Geological Survey, Reston, Va., 1984, \$5. *Water Resource Planning and Development*, M.S. Petersen, Prentice-Hall, Englewood Cliffs, N.J., xvii + 316 pp., 1984, \$35.

**Wandering Continents and Spreading Sea Floors on an Expanding Earth**, L. C. King, Wiley-Interscience, New York, x + 232 pp., 1983, \$29.95.

**Water Management Models in Practice: A Case Study of the Aswan High Dam**, D. Whittington and G. Guariso, Elsevier Scientific, New York, xvii + 246 pp., 1983, \$63.75.

**Water Resources: Distribution, Use, and Management**, J. R. Mather, J. Wiley & Sons, New York, xv + 439 pp., 1984, \$40.

**Water Wave Mechanics: For Engineers and Scientists**, R. G. Dean and R. A. Dalrymple, Prentice-Hall, Englewood Cliffs, N.J., xi + 353 pp., 1984, \$34.95.

**Waves on Fluid Interfaces**, R. E. Meyer (Ed.), Academic, New York, ix + 359 pp., 1983, \$25.

**Zeo-Agriculture: Use of Natural Zeolites in Agriculture and Aquaculture**, W. G. Pond and F. A. Mumpton (eds.), Westview, Boulder, xii + 296 pp., 1984, \$52.50.

**Yale University/Solid Earth Geophysics**. The Department of Geology and Geophysics is soliciting applications for a junior faculty position in solid earth geophysics to begin in the academic year 1985-1986. Areas of interest to the department include seismology, exploration geophysics, mechanical and physical properties of rocks and minerals, geomagnetism, tectonophysics, and geodesy. Curriculum vitae, publications and the names of three or more references should be sent by December 1, 1984 to: Dr. T. Turekian, Chairman, Department of Geology and Geophysics, Yale University, Box 6668, New Haven, CT 06511.

Yale University is an equal opportunity/affirmative action employer and encourages applications from all qualified scientists.

**Executive Director**. Executive Director of newly established Incorporated Research Institutions for Seismology (IRIS), a non-profit consortium of about 40 research universities. Initial duties include setting up Washington, D.C. office with associated financial and clerical services, conducting necessary contract negotiations with federal agencies and private organizations, handling procedures and arrangements for executive committee activities, and working with the managers of the various research programs. Under supervision of the President, represents the corporation as necessary in dealing with member institutions, funding agencies and contractors in administering large scientific programs. The corporation anticipates a level of research exceeding \$30 million annually in five years, with a permanent office staff of up to ten. Candidates must be able to work independently, with little staff support in the first year, and have sufficient breadth and experience to establish an efficient, functioning corporate office. Applicants should submit resumes and names of at least three references to: IRIS, Inc., Department ED, 2000 Florida Avenue, N.W., Washington, D.C. 20006.

IRIS is an equal opportunity employer.

**Applied Geophysics/Bowling Green State University**. The Department of Geology invites applications for a tenure track, assistant professor position in applied geophysics. Salary up to \$30,000; Ph.D. required. The successful candidate will be expected to develop a research program in some aspect of applied geophysics and teach courses in geophysics, exploration geophysics, and in his or her specialty.

The Department has 11 full-time faculty. In addition, two faculty from the Physics Department participate in our geophysics program. Complete physical instrumentation, including a seismograph station and rock mechanics lab, are available.

Interested persons should send resume, statement of research interests, official transcripts, and letters of reference to: Charles M. Onach, Chairman, Search Committee, Department of Geology, Bowling Green State University, Bowling Green, Ohio 43403. The closing date is November 1, 1984. We will be interviewing at BGSU in Reno, NV.

BGSU is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation of oceanic turbulence data. The tenure is for one or two years. The successful candidate should have a Ph.D. in physical oceanography and although experience with turbulence data is preferable it is not essential. The opportunity for involvement in data gathering expeditions is also available. Resumes can be sent to: Dr. R.G. Lueck, Code 68 LL, Naval Postgraduate School, Monterey, CA 93943. AA/EOE.

**Hydrogeologist/Illinois State Geological Survey**. Positions are available for research staff to study problems involving hazardous and radioactive waste disposal and groundwater resource evaluation. Applicants should possess an advanced degree in hydrogeology or related field and have strong communication skills. Send a letter of application, resume, list of three references and list of publications to: Marilyn-Rebecca, Personnel Office, Illinois State Geological Survey, 615 E. Peabody Drive, Champaign, IL 61820 by October 31, 1984.

The Illinois State Geological Survey is an equal opportunity/affirmative action employer.

**Postdoctoral Position/Naval Postgraduate School**. The Ocean Turbulence Laboratory has available a postdoctoral position for a person interested in the analysis and interpretation



## Dean College of Mathematical and Physical Sciences

The Ohio State University invites nominations and applications for the position of Dean of the College of Mathematical and Physical Sciences. The College consists of seven departments: Astronomy, Chemistry, Geodetic Science and Surveying, Geology and Mineralogy, Mathematics, Physics, and Statistics. Undergraduate, masters, and doctoral degree programs are offered in each department. Included in the College is an undergraduate program in computer and information science in cooperation with the Department of Computer and Information Science (College of Engineering), an interdisciplinary major in mathematical sciences, a major in actuarial science, and a major in Engineering Physics. The Materials Research Laboratory is an interdisciplinary program at the graduate level which involves Physics, Chemistry, Metallurgical Engineering, and Ceramic Engineering.

The College has a faculty of 230 FTE and teaches over 9,000 students (FTE) per year. Several of its departments are among the most distinguished in the University.

The Dean provides leadership as the chief executive of the College and promotes the goals of the College, departments and programs; the Dean reports to the Provost of the University. Qualifications for the position include a distinguished scholarly record in research and teaching plus demonstrated leadership and administrative ability. Candidates must qualify for a tenured appointment in one of the departments of the College at the level of Professor. The deanship is a twelve-month position.

Position available July 1, 1985, or at an earlier date. Salary is negotiable and commensurate with background and experience. Nominations or letters of application, curriculum vitae, and names, addresses, and telephone numbers of four references are required.

The Search Committee will begin to review dossiers on October 10 and will continue to receive nominations and applications until a candidate is selected.

Applications and nominations should be addressed to:

Chair, Search Committee for a Dean of the  
College of Mathematical and Physical Sciences  
The Ohio State University  
Office of Academic Affairs  
203 Bricker Hall  
Columbus, Ohio 43210

**Graduate Assistantships in Physics, Space Physics and Atmospheric Sciences.** Assistantships are available for graduate students seeking M.S. and Ph.D. degrees in Space Physics, Atmospheric Sciences or Physics at the University of Alaska. Research areas include both Experimental and Theoretical studies in Space Plasma Physics, Solar Physics, Computational Physics, Radio Physics, Atomic and Molecular Spectroscopy, Atmospheric Optics, Atmospheric Dynamics, Atmospheric Chemistry, Physical Meteorology and Climatology. This research is conducted through the Geophysical Institute. The stipend is \$12,000 to \$15,000 per year depending on credentials. Students with degrees in Physics, Atmospheric Sciences, Electrical or Mechanical Engineering are encouraged to apply. For more information, write to Professor J.R. Kan, Head, Department of Space Physics and Atmospheric Sciences, or Professor G. Sivjee, Head, Department of Physics, University of Alaska, Fairbanks, Alaska 99701 or call 907-474-7513.

**Research Associate in Marine Geochemistry/Texas A&M University.** Applications are invited for a position of Research Associate in the Department of Oceanography, Texas A&M University. Candidates should possess a Ph.D. and strong background in solution geochemistry or related area. Position will be for at least 1.5 years, to study acetate adsorption on mineral surfaces in the presence of organic matter. Starting salary \$18,000/year. Send letter of application, transcripts, resume and names and addresses of two references to: Dr. John W. Morse, Department of Oceanography, Texas A&M University, College Station, TX 77843.

**University of Colorado/Visting Faculty and Postdoctoral Fellowships for Research in Atmospheric and Climate Dynamics, Environmental and Atmospheric Chemistry, Environmental Biochemistry and Geochemistry.** One year awards offered by the Cooperative Institute for Research in Environmental Sciences (CIRES). CIRES is supported by NOAA and the University of Colorado. Awards may be made to senior scientists, including those on sabbatical leave, or to recent Ph.D. recipients. Stipends average \$25,000 for twelve months. The program is open to scientists of all countries. Selection of awardees is made in part on the likelihood of active interaction with CIRES scientists and other research groups in Boulder. Senior CIRES-affiliated scientists working in these fields include James Avery, Susan Avery, Ben Bailey, Roger Barry, John Birks, Ray Falk, Lang Farmer, Fred Fehsenfeld, William Hays, Murray Johnson, Bruce Kerk, Eric Kraus, Uwe Radok, Robert Sami, Robert Sievers, and Harold Walton. Applications for 1985-86 academic year awards should be sent to: Professor R. Sievers, Director of CIRES, Visitor Fellowship Program, Campus Box 449, University of Colorado, Boulder, CO 80509. Include a vitae, publications list, and a brief outline of the proposed research. In addition, applicants should submit three letters of recommendation. Junior applicants should submit undergraduate and graduate transcripts. First consideration will be given to applications received by December 1. Final application deadline is February 15.

The University of Colorado is an affirmative action/equal opportunity employer.

**Bureau of Mineral Resources, Australian Senior Research Scientist/Principal Research Scientist.** Applications are invited from suitably qualified men and women for the position of Senior Research Scientist (SRSP/Principal Research Scientist (PRS)). **BACKGROUND:** The ability to detect and identify nuclear explosions is an important element in achieving the Australian government's objective of seeking a Comprehensive Test Ban treaty which would ban all nuclear tests by all states in all environments. A National Monitoring Centre (NMC) and an International Data Centre (IDC) are being established to assist the international monitoring of nuclear explosions by seismological means. This group of ten scientific and technical staff will be established within the Bureau of Mineral Resources, Geology and Geophysics. The leader of the Group will be the head of the Earthquake Seismology Section within the Division of Geophysics.

The initial task of the Group Leader (SRSP/PRS) will be to arrange funding and carry out the monitoring and data analysis roles and to develop these into an operating system.

**DUTIES:** To direct the operations of the NMC and the IDC for the seismological monitoring of nuclear explosions.

**QUALIFICATIONS:** Admission to degree of Doctor of Philosophy or equivalent qualification together with considerable experience in seismological research and applied research in managing a scientific team, the science of monitoring nuclear explosions, seismological data processing, signal enhancement techniques, and digital data processing would be advantageous.

**SALARY:** Senior Research Scientist \$432,367—\$72,741; Principal Research Scientist \$438,688—\$73,041.

Classification will depend on the successful candidate's qualifications and experience in scientific research.

**CONDITIONS:** Conditions of service include superannuation, long service leave, four weeks annual leave and removal expenses to Canberra.

**APPLICATIONS:** Together with full personal and professional details and the names of at least three referees should be sent to:

The Director  
Bureau of Mineral Resources  
PO Box 378  
CANBERRA CITY A.C.T. 2601  
AUSTRALIA

By 26 October 1984.

**Civil Engineering.** The University of Notre Dame is seeking applications for a tenure track Assistant or Associate Professor position in its Department of Civil Engineering. Applicants should have an earned Ph.D. in Civil Engineering or an appropriate discipline. Rank is open. Applicants must have an active research program and be committed to excellence in teaching. Applicants should submit a curriculum vitae and arrange to have three letters of recommendation sent to: Lawrence Lundgren, Chairman, Department of Geological Sciences, University of Rochester, Rochester, New York 14627.

The University of Rochester is an equal opportunity/affirmative action employer.

**Manager, Research Computer Facility.** The University of Oklahoma is looking for a person to manage a recently purchased VAX 11/780 computer facility dedicated to research in the Geosciences. Hardware and software are provided for image processing, seismic reflection data processing, and graphical display of geological, geophysical and geophysical data.

In addition the 11780s has 8MB of CPU memory, the system includes an array processor, line tape drives, five disk drives, a laser printer, a 36" electrostatic plotter, and two high resolution graphics workstations with a digitizing table. The image processing hardware includes a Gould-Deane 198500 processor with 16 image memory planes, real time disk memory and three high resolution color monitors.

The person selected must have at least a BS degree in science, math, engineering or related field; two years programming experience including FORTRAN; educational or computing experience in solid earth geophysics or meteorology. Experience with the VAX VMS operating system as well as supervisory experience are desired.

Salary is negotiable. People interested in the position should send a resume, copies of academic transcripts, and the names, addresses and telephone numbers of three references to:

John Wickham, Director  
School of Geology & Geophysics  
University of Oklahoma  
Norman, OK 73019

Applications must be received by November 2, 1984.

**University of Wisconsin—Madison.** The Department of Geology and Geophysics invites applications for an equipped tenure track position at the assistant professor level in applied geomorphology and/or hydrogeology commencing in August 1985. The applicant should be committed to developing a strong research program as well as teaching undergraduate courses in some aspects of geomorphology and environmental geology. The Ph.D. is required. Applicants with course work in engineering and an interest in the field application of geologic principles are especially encouraged to apply. Send letter of application outlining your professional goals, transcripts, resume, copies of publications, and three letters of reference to: Dr. Mary P. Anderson, Department of Geology and Geophysics, Weeks Hall, University of Wisconsin, Madison, WI 53706. Closing date is January 1, 1985.

The University of Wisconsin is an equal opportunity/affirmative action employer.

**Physical Scientist/Department of Commerce, National Oceanic and Atmospheric Administration (NOAA).** The National Environmental Satellite, Data, and Information Service (NESDIS), Office of Research and Applications (ORA) announces a vacancy for the position of Physical Scientist, GS-130-13, Suitland, Maryland. Vacancy closes October 31, 1984. The successful applicant will perform basic research and applied research in physical sciences with emphasis on the land sciences using imagery and physical measurements collected by meteorological and land resource satellites. The purpose of these investigations is to increase knowledge and understanding of hydrologic, climatic, and land-related phenomena and processes, and to develop applications of remote sensing data to hydrologic, climatic, and land-related problems. The position requires a demonstrated ability in scientific research on the application of remote sensing to the above stated problems, as evidenced by publications in the scientific literature. A Ph.D. in the physical sciences or equivalent, familiarity with programming of minicomputer computers and experience with interactive digital systems are desirable. Persons interested in applying must request a copy of the vacancy announcement which contains qualification requirements, by writing to NOAA, P.O. Box 308, Washington, D.C. 20253. ATTN: RAS/DC24, Mary Queen, or calling 301-769-1986. Applications should be submitted on Standard Form 171.

Department of Commerce is an equal opportunity employer. U.S. citizenship required.

**University of Rochester/Faculty Position.** The Department of Geological Sciences at the University of Rochester invites applications for a tenure track position in sedimentology. Rank is open. Applicants must have an active research program and be committed to excellence in teaching. Applicants should submit a curriculum vitae and arrange to have three letters of recommendation sent to: Lawrence Lundgren, Chairman, Department of Geological Sciences, University of Rochester, Rochester, New York 14627.

The University of Rochester is an equal opportunity/affirmative action employer.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days, April 16-18, 1985, Fort Collins, Colorado.** Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

magnetic, magnetic, paleomagnetic, gravimetric, and geothermic techniques; seismic activity and earthquakes; magnetism, including extrusion and intrusion, formation of ore bodies and kimberlites, and formation of oil reservoirs; petrological, geomorphological, and geochronometric investigations; and rheology of rocks and geodynamic modeling. There will also be a tour of the Panxi Rifting Belt, and a second tour in Northern China is being planned. The official languages of the symposium will be Chinese and English.

**Future AGU Meetings**

**Fall Meetings**  
Dec. 3-7, 1984, San Francisco, California. Call for papers appeared in July 3, 1984 Eos.

Dec. 9-13, 1985, San Francisco, California. Abstracts due mid-September 1985.

Dec. 8-12, 1986, San Francisco, California.

**Spring Meetings**  
May 27-31, 1985, Baltimore, Maryland. Abstracts due early March 1985.

May 19-23, 1986, Baltimore, Maryland.

**Regional Meetings**  
**Front Range Branch Symposium on Geophysics and Geology of Yellowstone,** October 25, 1984, Golden, Colorado. Abstracts due October 12, 1984; call for papers appeared in September 18, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**Front Range Branch Hydrology Days,** April 16-18, 1985, Fort Collins, Colorado. Abstracts due December 31, 1984 for professional hydrologists. February 15, 1985 for students; call for papers appeared in July 24, 1984 Eos.

**AGU Separates**

To Order: The order number can be found at the end of each abstract; use all digits when ordering. Only papers with order numbers are available from AGU. Cost: \$3.50 for the first article and \$1.00 for each additional article in the same order. Payment must accompany order. Deposit accounts available.

Send your order to:  
American Geophysical Union  
2000 Florida Avenue, N.W.  
Washington, D.C. 20009

**Aeronomy**

**0430 Composition (Molecular)**  
AN UPPER LIMIT FOR HYDROGENATED HYDROGEN PERIODS  
R. V. CHANE (Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138) and W. A. TRACH

Measurements of an altitude-dependent upper limit for hydrogenated hydrogen peroxide have been obtained from the stratosphere with a thermal emission sensing spectrometer during a balloon flight on 23 October 1983. The 20 ppb limit reaches a minimum of 0.5 ppb near 26 km, increases to 1.5 ppb near 30 km, and then decreases to 0.5 ppb near 35 km. This upper limit is compared with two theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions.

**0430 Composition (Molecular)**  
AN UPPER LIMIT FOR HYDROGENATED HYDROGEN PERIODS  
R. V. CHANE (Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138) and W. A. TRACH

Measurements of an altitude-dependent upper limit for hydrogenated hydrogen peroxide have been obtained from the stratosphere with a thermal emission sensing spectrometer during a balloon flight on 23 October 1983. The 20 ppb limit reaches a minimum of 0.5 ppb near 26 km, increases to 1.5 ppb near 30 km, and then decreases to 0.5 ppb near 35 km. This upper limit is compared with two theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions.

**0430 Composition (Molecular)**  
AN UPPER LIMIT FOR HYDROGENATED HYDROGEN PERIODS  
R. V. CHANE (Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138) and W. A. TRACH

Measurements of an altitude-dependent upper limit for hydrogenated hydrogen peroxide have been obtained from the stratosphere with a thermal emission sensing spectrometer during a balloon flight on 23 October 1983. The 20 ppb limit reaches a minimum of 0.5 ppb near 26 km, increases to 1.5 ppb near 30 km, and then decreases to 0.5 ppb near 35 km. This upper limit is compared with two theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions.

**0430 Composition (Molecular)**  
AN UPPER LIMIT FOR HYDROGENATED HYDROGEN PERIODS  
R. V. CHANE (Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138) and W. A. TRACH

Measurements of an altitude-dependent upper limit for hydrogenated hydrogen peroxide have been obtained from the stratosphere with a thermal emission sensing spectrometer during a balloon flight on 23 October 1983. The 20 ppb limit reaches a minimum of 0.5 ppb near 26 km, increases to 1.5 ppb near 30 km, and then decreases to 0.5 ppb near 35 km. This upper limit is compared with two theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions.

**0430 Composition (Molecular)**  
AN UPPER LIMIT FOR HYDROGENATED HYDROGEN PERIODS  
R. V. CHANE (Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138) and W. A. TRACH

Measurements of an altitude-dependent upper limit for hydrogenated hydrogen peroxide have been obtained from the stratosphere with a thermal emission sensing spectrometer during a balloon flight on 23 October 1983. The 20 ppb limit reaches a minimum of 0.5 ppb near 26 km, increases to 1.5 ppb near 30 km, and then decreases to 0.5 ppb near 35 km. This upper limit is compared with two theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions.

**0430 Composition (Molecular)**  
AN UPPER LIMIT FOR HYDROGENATED HYDROGEN PERIODS  
R. V. CHANE (Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138) and W. A. TRACH

Measurements of an altitude-dependent upper limit for hydrogenated hydrogen peroxide have been obtained from the stratosphere with a thermal emission sensing spectrometer during a balloon flight on 23 October 1983. The 20 ppb limit reaches a minimum of 0.5 ppb near 26 km, increases to 1.5 ppb near 30 km, and then decreases to 0.5 ppb near 35 km. This upper limit is compared with two theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions.

**0430 Composition (Molecular)**  
AN UPPER LIMIT FOR HYDROGENATED HYDROGEN PERIODS  
R. V. CHANE (Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138) and W. A. TRACH

Measurements of an altitude-dependent upper limit for hydrogenated hydrogen peroxide have been obtained from the stratosphere with a thermal emission sensing spectrometer during a balloon flight on 23 October 1983. The 20 ppb limit reaches a minimum of 0.5 ppb near 26 km, increases to 1.5 ppb near 30 km, and then decreases to 0.5 ppb near 35 km. This upper limit is compared with two theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions.

**0430 Composition (Molecular)**  
AN UPPER LIMIT FOR HYDROGENATED HYDROGEN PERIODS  
R. V. CHANE (Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138) and W. A. TRACH

Measurements of an altitude-dependent upper limit for hydrogenated hydrogen peroxide have been obtained from the stratosphere with a thermal emission sensing spectrometer during a balloon flight on 23 October 1983. The 20 ppb limit reaches a minimum of 0.5 ppb near 26 km, increases to 1.5 ppb near 30 km, and then decreases to 0.5 ppb near 35 km. This upper limit is compared with two theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions.

**0430 Composition (Molecular)**  
AN UPPER LIMIT FOR HYDROGENATED HYDROGEN PERIODS  
R. V. CHANE (Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138) and W. A. TRACH

Measurements of an altitude-dependent upper limit for hydrogenated hydrogen peroxide have been obtained from the stratosphere with a thermal emission sensing spectrometer during a balloon flight on 23 October 1983. The 20 ppb limit reaches a minimum of 0.5 ppb near 26 km, increases to 1.5 ppb near 30 km, and then decreases to 0.5 ppb near 35 km. This upper limit is compared with two theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions.

**0430 Composition (Molecular)**  
AN UPPER LIMIT FOR HYDROGENATED HYDROGEN PERIODS  
R. V. CHANE (Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138) and W. A. TRACH

Measurements of an altitude-dependent upper limit for hydrogenated hydrogen peroxide have been obtained from the stratosphere with a thermal emission sensing spectrometer during a balloon flight on 23 October 1983. The 20 ppb limit reaches a minimum of 0.5 ppb near 26 km, increases to 1.5 ppb near 30 km, and then decreases to 0.5 ppb near 35 km. This upper limit is compared with two theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5 to 1.5 ppb, which is slightly higher than the theoretical predictions. The upper limit for hydrogenated hydrogen peroxide is in the range of 0.5







